

t50_classes1
(TMFJ7Ub2LdDpNKpky4jLPP9XWjCaJDy4C4w)

October 27, 2020

Let $v1_ordinal1 : \iota \Rightarrow o$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1_tarski X0 X1) \wedge (r1_tarski X0 X2)) \Rightarrow (r1_tarski X0 (k3_xboole_0 X1 X2)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k3_xboole_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (X3 \in X1))) \quad (2)$$

Assume the following.

$$\forall X0. (v1_ordinal1 X0) \Leftrightarrow (\forall X1. (X1 \in X0) \Rightarrow (r1_tarski X1 X0)) \quad (3)$$

Theorem 1

$$\forall X0. \forall X1. ((v1_ordinal1 X0) \wedge (v1_ordinal1 X1)) \Rightarrow (v1_ordinal1 (k3_xboole_0 X0 X1))$$