

t50\_cohsp\_1  
(TMdrrK25AkV9p1fSjTevsGsvqcMKa1nv4gV)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $k4\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k10\_cohsp\_1 : \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $k7\_cohsp\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. (k4\_tarski X0 X1 = k4\_tarski X2 X3) \Rightarrow ((X0 = X2) \wedge (X1 = X3)) \quad (1)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1. (X1 = k10\_cohsp\_1 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. \exists X4. (X2 = k4\_tarski X3 X4) \wedge (k4\_tarski (k1\_tarski X3) X4 \in k7\_cohsp\_1 X0)))))) \quad (2)$$

**Theorem 1**

$$\forall X0. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1. \forall X2. (k4\_tarski X1 X2 \in k10\_cohsp\_1 X0) \Leftrightarrow (k4\_tarski (k1\_tarski X1) X2 \in k7\_cohsp\_1 X0))$$