

t50_ordinal6 (TM-
ceF5JJxG6BJrMUfPPDsLUx1wSwCqx6HYv)

October 27, 2020

Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v5_ordinal1 : \iota \Rightarrow o$ be given. Let $v3_ordinal6 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $r1_abian : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_ordinal6 : \iota \Rightarrow \iota$ be given. Let $v1_ordinal6 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $v1_ordinal2 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_ordinal6 : \iota \Rightarrow \iota$ be given. Let $k3_wellord1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_wellord2 : \iota \Rightarrow \iota$ be given. Let $k1_ordinal6 : \iota \Rightarrow \iota$ be given. Let $k2_ordinal1 : \iota \Rightarrow \iota$ be given. Let $k2_wellord2 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v3_ordinal1 X0) \Rightarrow ((X0 \neq k1_xboole_0) \Rightarrow (k1_xboole_0 \in X0)) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v5_ordinal1 \\ & X0) \wedge (v3_ordinal6 X0)))) \Rightarrow (v1_ordinal6 (ReplSep (toset (\lambda X1 : \\ & \iota.m1_subset_1 X1 (k9_xtuple_0 (k1_funct_1 X0 k6_numbers)))) \\ & (\lambda X1 : \iota.(X1 \in k9_xtuple_0 (k1_funct_1 X0 k6_numbers)) \wedge (\forall X2. \\ & ((v1_relat_1 X2) \wedge ((v1_funct_1 X2) \wedge ((v5_ordinal1 X2) \wedge (v1_ordinal2 \\ & X2)))) \Rightarrow ((X2 \in k10_xtuple_0 X0) \Rightarrow (r1_abian X1 X2)))) (\lambda X1 : \iota. \\ & X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (3)$$

Assume the following.

$$\forall X0.(v1_ordinal6 X0) \Rightarrow (k10_xtuple_0 (k2_ordinal6 X0) = X0) \quad (4)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v5_ordinal1 X0))) \Rightarrow (v3_ordinal1 (k9_xtuple_0 X0)) \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v3_ordinal6 X0))) \Rightarrow ((v1_relat_1 (k1_funct_1 X0 X1)) \wedge (v1_funct_1 (k1_funct_1 X0 X1))) \quad (7)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v5_ordinal1 X0) \wedge (v3_ordinal6 X0)))) \Rightarrow ((v1_relat_1 (k7_ordinal6 X0)) \wedge ((v1_funct_1 (k7_ordinal6 X0)) \wedge ((v5_ordinal1 (k7_ordinal6 X0)) \wedge (v1_ordinal2 (k7_ordinal6 X0))))) \quad (8)$$

Assume the following.

$$\forall X0. k2_ordinal6 X0 = k3_wellord1 (k1_wellord2 (k1_ordinal6 X0)) (k1_wellord2 (k2_ordinal1 X0)) \quad (9)$$

Assume the following.

$$\forall X0. k1_ordinal6 X0 = k2_wellord2 (k1_wellord2 (k2_ordinal1 X0)) \quad (10)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. (X1 = k10_xtuple_0 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. (X3 \in k9_xtuple_0 X0) \wedge (X2 = k1_funct_1 X0 X3)))) \quad (11)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_abian X0 X1) \Leftrightarrow ((X0 \in k9_xtuple_0 X1) \wedge (X0 = k1_funct_1 X1 X0))) \quad (12)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v5_ordinal1 X0) \wedge (v3_ordinal6 X0)))) \Rightarrow (k7_ordinal6 X0 = k2_ordinal6 (ReplSep (toset (\lambda X1 : \iota. m1_subset_1 X1 (k9_xtuple_0 (k1_funct_1 X0 k6_numbers)))) (\lambda X1 : \iota. (X1 \in k9_xtuple_0 (k1_funct_1 X0 k6_numbers)))) \wedge (\forall X2. ((v1_relat_1 X2) \wedge ((v1_funct_1 X2) \wedge ((v5_ordinal1 X2) \wedge (v1_ordinal2 X2)))) \Rightarrow ((X2 \in k10_xtuple_0 X0) \Rightarrow (r1_abian X1 X2)))) (\lambda X1 : \iota. X1)) \quad (13)$$

Assume the following.

$$\forall X0.(v1_ordinal6\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ X0)\Rightarrow(v3_ordinal1\ X1)) \quad (14)$$

Assume the following.

$$\forall X0.(v3_ordinal1\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ X0)\Rightarrow(v3_ordinal1\ X1)) \quad (15)$$

Assume the following.

$$\forall X0.(v3_ordinal1\ X0)\Rightarrow(v1_ordinal6\ X0) \quad (16)$$

Theorem 1

$$\begin{aligned} &\forall X0.(v3_ordinal1\ X0)\Rightarrow(\forall X1.((v1_relat_1\ X1)\wedge((\\ &v1_funct_1\ X1)\wedge((v5_ordinal1\ X1)\wedge(v3_ordinal6\ X1))))\Rightarrow(\neg(k9_xtuple_0 \\ &X1\neq k1_xboole_0)\wedge((\forall X2.(v3_ordinal1\ X2)\Rightarrow((X2\in k9_xtuple_0 \\ &X1)\Rightarrow(r1_abian\ X0\ (k1_funct_1\ X1\ X2))))\wedge(\forall X2.(v3_ordinal1 \\ &X2)\Rightarrow(\neg(X2\in k9_xtuple_0\ (k7_ordinal6\ X1))\wedge(X0 = k1_funct_1\ (k7_ordinal6 \\ &X1\ X2)))))) \end{aligned}$$