

t50_qc_lang3
(TMMu7TpKxYt6UcCTqrauTwsn1vPGJuXbFJZ)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k4_qc_lang3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_qc_lang2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_qc_lang2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k13_qc_lang2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k14_qc_lang2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k19_qc_lang1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_qc_lang2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k20_qc_lang1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0) \Rightarrow (\forall X2.((\neg v1_xboole_0 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_qc_lang1 X0)))) \Rightarrow ((v3_qc_lang2 X1 X0) \Rightarrow (k4_qc_lang3 X0 X2 X1 = \\ k4_subset_1 X2 (k4_qc_lang3 X0 X2 (k13_qc_lang2 X0 X1)) (k4_qc_lang3 \\ X0 X2 (k14_qc_lang2 X0 X1))))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0) \Rightarrow (\forall X2.(m1_subset_1 X2 (k9_qc_lang1 X0) \Rightarrow ((k13_qc_lang2 \\ X0 (k4_qc_lang2 X0 X1 X2) = X1) \wedge ((k14_qc_lang2 X0 (k4_qc_lang2 X0 \\ X1 X2) = X2) \wedge ((k19_qc_lang1 X0 (k4_qc_lang2 X0 X1 X2) = k2_qc_lang2 \\ X0 X1 X2) \wedge (k20_qc_lang1 X0 (k4_qc_lang2 X0 X1 X2) = k2_qc_lang2 X0 \\ X2 X1))))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((m1_qc_lang1 X0) \wedge ((m1_subset_1 \\ X1 (k9_qc_lang1 X0) \wedge (m1_subset_1 X2 (k9_qc_lang1 X0)))) \Rightarrow (m1_subset_1 \\ (k4_qc_lang2 X0 X1 X2) (k9_qc_lang1 X0))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k9_qc_lang1 \\ X0)) \Rightarrow ((v3_qc_lang2\ X1\ X0) \Leftrightarrow (\exists X2.(m1_subset_1\ X2\ (k9_qc_lang1 \\ X0)) \wedge (\exists X3.(m1_subset_1\ X3\ (k9_qc_lang1\ X0)) \wedge (X1 = k4_qc_lang2 \\ X0\ X2\ X3)))))) \end{aligned} \quad (4)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k9_qc_lang1 \\ X0)) \Rightarrow (\forall X2.(m1_subset_1\ X2\ (k9_qc_lang1\ X0)) \Rightarrow (\forall X3. \\ ((\neg v1_xboole_0\ X3) \wedge (m1_subset_1\ X3\ (k1_zfmisc_1\ (k2_qc_lang1 \\ X0)))) \Rightarrow (k4_qc_lang3\ X0\ X3\ (k4_qc_lang2\ X0\ X1\ X2) = k4_subset_1\ X3 \\ (k4_qc_lang3\ X0\ X3\ X1)\ (k4_qc_lang3\ X0\ X3\ X2)))))) \end{aligned}$$