

t50_tdlat_2 (TMc- neiP3ngwzeJHogMzAaymGtQ6VMEDAohp)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_pre_topc : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tops_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_setfam_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_pcomps_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tdlat_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v2_pre_topc X0) \wedge (l1_pre_topc X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 (u1_struct_0 X0)))) \Rightarrow \\ & (r1_tarski (k1_tops_1 X0 (k6_setfam_1 (u1_struct_0 X0) X1)) (k6_setfam_1 \\ & (u1_struct_0 X0) (k1_tdlat_2 X0 X1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1_tarski X0 X1) \wedge (r1_tarski X1 X2)) \Rightarrow (r1_tarski X0 X2) \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. (l1_pre_topc X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 \\ & (u1_struct_0 X0))) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ & (u1_struct_0 X0))) \Rightarrow ((r1_tarski X1 X2) \Rightarrow (r1_tarski (k2_pre_topc \\ & X0 X1) (k2_pre_topc X0 X2)))))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v2_pre_topc X0) \wedge (l1_pre_topc X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 (u1_struct_0 X0)))) \Rightarrow \\ & (r1_tarski (k2_pre_topc X0 (k6_setfam_1 (u1_struct_0 X0) X1)) \\ & (k6_setfam_1 (u1_struct_0 X0) (k1_pcomps_1 X0 X1)))) \end{aligned} \tag{4}$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 X0))) \Rightarrow (m1_subset_1 (k6_setfam_1 X0 X1) (k1_zfmisc_1 X0)) \tag{5}$$

Assume the following.

$$\forall X0.\forall X1.((l1_pre_topc\ X0)\wedge(m1_subset_1\ X1\ (k1_zfmisc_1\ (u1_struct_0\ X0))))\Rightarrow(m1_subset_1\ (k1_tops_1\ X0\ X1)\ (k1_zfmisc_1\ (u1_struct_0\ X0))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(((v2_pre_topc\ X0)\wedge(l1_pre_topc\ X0))\wedge(m1_subset_1\ X1\ (k1_zfmisc_1\ (k1_zfmisc_1\ (u1_struct_0\ X0)))))\Rightarrow(m1_subset_1\ (k1_tdlat_2\ X0\ X1)\ (k1_zfmisc_1\ (k1_zfmisc_1\ (u1_struct_0\ X0)))) \quad (7)$$

Theorem 1

$$\forall X0.((\neg v2_struct_0\ X0)\wedge((v2_pre_topc\ X0)\wedge(l1_pre_topc\ X0)))\Rightarrow(\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1\ (k1_zfmisc_1\ (u1_struct_0\ X0))))\Rightarrow(r1_tarski\ (k2_pre_topc\ X0\ (k1_tops_1\ X0\ (k6_setfam_1\ (u1_struct_0\ X0)\ X1)))\ (k6_setfam_1\ (u1_struct_0\ X0)\ (k1_pcomps_1\ X0\ (k1_tdlat_2\ X0\ X1))))))$$