

t51\_bvfunc26 (TMX-  
pXi8vECquyQgw3hPnZEn5pvRCsDbMTYE)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_margrel1 : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r2\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_bvfunc\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_bvfunc26 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_bvfunc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_bvfunc\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned}
 & \forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. ((v1\_funct\_1 X1) \wedge ( \\
 & (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\
 & k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X2. ((v1\_funct\_1 X2) \wedge \\
 & ((v1\_funct\_2 X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\
 & (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X3. ((v1\_funct\_1 X3) \wedge \\
 & ((v1\_funct\_2 X3 X0 k6\_margrel1) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 \\
 & (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (r2\_funct\_2 X0 k6\_margrel1 \\
 & (k5\_bvfunc\_1 X0 (k5\_bvfunc\_1 X0 X1 X2) X3) (k5\_bvfunc\_1 X0 X1 (k5\_bvfunc\_1 \\
 & X0 X2 X3))))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
 & \forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. ((v1\_funct\_1 X1) \wedge ( \\
 & (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\
 & k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X2. ((v1\_funct\_1 X2) \wedge \\
 & ((v1\_funct\_2 X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\
 & (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (r2\_funct\_2 X0 k6\_margrel1 \\
 & (k5\_bvfunc26 X0 X1 X2) (k1\_bvfunc\_1 X0 (k2\_bvfunc\_1 X0 X1 X2))))))
 \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned}
& \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((v1\_funct\_1 X1) \wedge \\
& (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\
& k2\_zfmisc\_1 X0 k6\_margrel1)))) \Rightarrow (\forall X2.((v1\_funct\_1 X2) \wedge \\
& (v1\_funct\_2 X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\
& (k2\_zfmisc\_1 X0 k6\_margrel1)))) \Rightarrow (r2\_funct\_2 X0 k6\_margrel1 \\
& (k1\_bvfunc\_1 X0 (k2\_bvfunc\_1 X0 X1 X2)) (k5\_bvfunc\_1 X0 (k1\_bvfunc\_1 \\
& X0 X1) (k1\_bvfunc\_1 X0 X2))))))
\end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned}
& \forall X0.\forall X1.\forall X2.\forall X3.(((v1\_funct\_1 X2) \wedge \\
& ((v1\_funct\_2 X2 X0 X1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\
& X0 X1)))) \wedge ((v1\_funct\_1 X3) \wedge ((v1\_funct\_2 X3 X0 X1) \wedge (m1\_subset\_1 \\
& X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))))) \Rightarrow ((r2\_funct\_2 X0 X1 X2 \\
& X3) \Leftrightarrow (X2 = X3))
\end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned}
& \forall X0.\forall X1.\forall X2.((\neg v1\_xboole\_0 X0) \wedge (((v1\_funct\_1 \\
& X1) \wedge ((v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 \\
& (k2\_zfmisc\_1 X0 k6\_margrel1)))) \wedge ((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 \\
& X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\
& X0 k6\_margrel1)))))) \Rightarrow ((v1\_funct\_1 (k5\_bvfunc\_1 X0 X1 X2)) \wedge ( \\
& (v1\_funct\_2 (k5\_bvfunc\_1 X0 X1 X2) X0 k6\_margrel1) \wedge (m1\_subset\_1 \\
& (k5\_bvfunc\_1 X0 X1 X2) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1))))))
\end{aligned} \tag{5}$$

Assume the following.

$$\begin{aligned}
& \forall X0.\forall X1.\forall X2.((\neg v1\_xboole\_0 X0) \wedge (((v1\_funct\_1 \\
& X1) \wedge ((v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 \\
& (k2\_zfmisc\_1 X0 k6\_margrel1)))) \wedge ((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 \\
& X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\
& X0 k6\_margrel1)))))) \Rightarrow ((v1\_funct\_1 (k5\_bvfunc26 X0 X1 X2)) \wedge ( \\
& (v1\_funct\_2 (k5\_bvfunc26 X0 X1 X2) X0 k6\_margrel1) \wedge (m1\_subset\_1 \\
& (k5\_bvfunc26 X0 X1 X2) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1))))))
\end{aligned} \tag{6}$$

Assume the following.

$$\begin{aligned}
& \forall X0.\forall X1.\forall X2.((\neg v1\_xboole\_0 X0) \wedge (((v1\_funct\_1 \\
& X1) \wedge ((v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 \\
& (k2\_zfmisc\_1 X0 k6\_margrel1)))) \wedge ((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 \\
& X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\
& X0 k6\_margrel1)))))) \Rightarrow ((v1\_funct\_1 (k2\_bvfunc\_1 X0 X1 X2)) \wedge ( \\
& (v1\_funct\_2 (k2\_bvfunc\_1 X0 X1 X2) X0 k6\_margrel1) \wedge (m1\_subset\_1 \\
& (k2\_bvfunc\_1 X0 X1 X2) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1))))))
\end{aligned} \tag{7}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge ((v1\_funct\_1 X1) \wedge \\ & (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\ & k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow ((v1\_funct\_1 (k1\_bvfunc\_1 \\ X0 X1)) \wedge ((v1\_funct\_2 (k1\_bvfunc\_1 X0 X1) X0 k6\_margrel1) \wedge (m1\_subset\_1 \\ (k1\_bvfunc\_1 X0 X1) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((\neg v1\_xboole\_0 X0) \wedge (((v1\_funct\_1 \\ X1) \wedge ((v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 \\ (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \wedge ((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 \\ X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ X0 k6\_margrel1)))))) \Rightarrow (k5\_bvfunc\_1 X0 X1 X2 = k5\_bvfunc\_1 X0 X2 \\ X1) \end{aligned} \quad (9)$$

**Theorem 1**

$$\begin{aligned} & \forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. ((v1\_funct\_1 X1) \wedge \\ & (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\ & k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X2. ((v1\_funct\_1 X2) \wedge \\ & ((v1\_funct\_2 X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X3. ((v1\_funct\_1 X3) \wedge \\ & ((v1\_funct\_2 X3 X0 k6\_margrel1) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (r2\_funct\_2 X0 k6\_margrel1 \\ & (k5\_bvfunc\_1 X0 X1 (k5\_bvfunc26 X0 X2 X3)) (k5\_bvfunc\_1 X0 (k5\_bvfunc\_1 \\ & X0 X1 (k1\_bvfunc\_1 X0 X2)) (k1\_bvfunc\_1 X0 X3)))))) \end{aligned}$$