

t51_card_2

(TMKvqM99U1BN4kQKnCxyFtYoKcXR5SR7Y4z)

October 27, 2020

Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_card_1 : \iota \Rightarrow \iota$ be given. Let $k1_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_3 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (\forall X3.(v1_xreal_0 X3) \Rightarrow (((r1_xxreal_0 \\ & X0 X1) \wedge (r1_xxreal_0 X2 X3)) \Rightarrow (r1_xxreal_0 (k2_xcmplx_0 X0 X2) (\\ & k2_xcmplx_0 X1 X3)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. r1_xxreal_0 (k5_card_1 (k2_tarski X0 X1)) \quad np_2 \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_finset_1 X0) \Rightarrow (\forall X1.(v1_finset_1 X1) \Rightarrow (r1_xxreal_0 \\ & (k5_card_1 (k2_xboole_0 X0 X1)) (k2_nat_1 (k5_card_1 X0) (k5_card_1 \\ & X1)))) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0. k1_card_1 (k1_tarski X0) = np_1 \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow \\ & (r1_xxreal_0 X0 X2)))) \end{aligned} \tag{5}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.k1_enumset1\ X0\ X1\ X2 = k2_xboole_0\ (k1_tarski\ X0)\ (k2_tarski\ X1\ X2) \quad (6)$$

Assume the following.

$$((v2_xxreal_0\ np_3)\wedge(m2_subset_1\ np_3\ k1_numbers\ k5_numbers))\wedge\ ((m1_subset_1\ np_3\ k5_numbers)\wedge(m1_subset_1\ np_3\ k1_numbers)) \quad (7)$$

Assume the following.

$$((v2_xxreal_0\ np_2)\wedge(m2_subset_1\ np_2\ k1_numbers\ k5_numbers))\wedge\ ((m1_subset_1\ np_2\ k5_numbers)\wedge(m1_subset_1\ np_2\ k1_numbers)) \quad (8)$$

Assume the following.

$$((v2_xxreal_0\ np_1)\wedge(m2_subset_1\ np_1\ k1_numbers\ k5_numbers))\wedge\ ((m1_subset_1\ np_1\ k5_numbers)\wedge(m1_subset_1\ np_1\ k1_numbers)) \quad (9)$$

Assume the following.

$$k2_xcmplx_0\ np_1\ np_2 = np_3 \quad (10)$$

Assume the following.

$$r1_xxreal_0\ np_1\ np_1 \quad (11)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (12)$$

Assume the following.

$$\forall X0.(v1_finset_1\ X0)\Rightarrow(k5_card_1\ X0 = k1_card_1\ X0) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1\ X0\ k5_numbers)\wedge(v7_ordinal1\ X1))\Rightarrow(k2_nat_1\ X0\ X1 = k2_xcmplx_0\ X0\ X1) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0\ X0)\wedge(v1_xreal_0\ X1))\Rightarrow(v1_xreal_0\ (k2_xcmplx_0\ X0\ X1)) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.v1_finset_1\ (k1_enumset1\ X0\ X1\ X2) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.v1_finset_1\ (k2_tarski\ X0\ X1) \quad (17)$$

Assume the following.

$$\forall X0.v1_finset_1 (k1_tarski X0) \quad (18)$$

Assume the following.

$$\forall X0.(v1_finset_1 X0) \Rightarrow (m1_subset_1 (k5_card_1 X0) k4_ordinal1) \quad (19)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (20)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (21)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_xreal_0 X0) \quad (22)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (23)$$

Theorem 1

$$\forall X0.\forall X1.\forall X2.r1_xxreal_0 (k5_card_1 (k1_enumset1 X0 X1 X2)) \quad np_3$$