

t51_card_3 (TM- FrioRRsEb1gTPdpfCuyuYVMRCKUGQofRC)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_card_3 : \iota \Rightarrow \iota$ be given. Let $k8_card_3 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. r1_tarski X0 X0 \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. (X1 = \\ k4_card_3 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. ((v1_relat_1 \\ X3) \wedge (v1_funct_1 X3)) \wedge ((X2 = X3) \wedge ((r1_tarski (k9_xtuple_0 X3) \\ (k9_xtuple_0 X0)) \wedge (\forall X4. (X4 \in k9_xtuple_0 X3) \Rightarrow (k1_funct_1 \\ X3 X4 \in k1_funct_1 X0 X4))))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. (X1 = \\ k4_card_3 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. ((v1_relat_1 \\ X3) \wedge (v1_funct_1 X3)) \wedge ((X2 = X3) \wedge ((k9_xtuple_0 X3 = k9_xtuple_0 \\ X0) \wedge (\forall X4. (X4 \in k9_xtuple_0 X0) \Rightarrow (k1_funct_1 X3 X4 \in k1_funct_1 \\ X0 X4))))))) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \tag{4}$$

Theorem 1

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (r1_tarski (k4_card_3 X0) (k8_card_3 X0))$$