

t51\_fomodel0

(TMKc1Dt2P7K6zoF2g1H4wffFBWSuYCnTUdL)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.(( \\ v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_tarski X0 X1) \Leftrightarrow ((r1\_tarski \\ (k9\_xtuple\_0 X0) (k9\_xtuple\_0 X1)) \wedge (\forall X2.(X2 \in k9\_xtuple\_0 \\ X0) \Rightarrow (k1\_funct\_1 X0 X2 = k1\_funct\_1 X1 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (r1\_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (2)$$

**Theorem 1**

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.(( \\ v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_tarski X0 X1) \Leftrightarrow (\forall X2. \\ (X2 \in k9\_xtuple\_0 X0) \Rightarrow ((X2 \in k9\_xtuple\_0 X1) \wedge (k1\_funct\_1 X0 X2 = \\ k1\_funct\_1 X1 X2)))))) \end{aligned}$$