

# t51\_ordinal3

## (TMT9PfuJdSPPERboLhugckv67G8EMKL6LZ)

October 27, 2020

Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k10\_ordinal2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_ordinal3 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_ordinal1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Assume the following.

$$\forall X0.(v3\_ordinal1 X0) \Rightarrow (\forall X1.(v3\_ordinal1 X1) \Rightarrow ((r1\_ordinal1 X0 X1) \vee (X1 \in X0))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((v3\_ordinal1 X0) \wedge (v3\_ordinal1 X1)) \Rightarrow (v3\_ordinal1 (k5\_ordinal3 X0 X1)) \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v3\_ordinal1 X0) \Rightarrow (\forall X1.(v3\_ordinal1 X1) \Rightarrow (\forall X2. \\ & (v3\_ordinal1 X2) \Rightarrow (((r1\_ordinal1 X1 X0) \Rightarrow ((X2 = k5\_ordinal3 X0 X1) \Leftrightarrow \\ & (X0 = k10\_ordinal2 X1 X2))) \wedge ((\neg r1\_ordinal1 X1 X0) \Rightarrow ((X2 = k5\_ordinal3 \\ & X0 X1) \Leftrightarrow (X2 = k1\_xboole\_0)))))) \quad (3) \end{aligned}$$

Assume the following.

$$\forall X0.\forall X1.(X0 \in X1) \Rightarrow (\neg X1 \in X0) \quad (4)$$

### Theorem 1

$$\forall X0.(v3\_ordinal1 X0) \Rightarrow (\forall X1.(v3\_ordinal1 X1) \Rightarrow ((X0 \in X1) \Rightarrow (X1 = k10\_ordinal2 X0 (k5\_ordinal3 X1 X0))))$$