

t51_qc_lang3
(TMZCWFWeR8uGrzfqTFiT7syb5THo4H5mCyk)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $v4_qc_lang2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_qc_lang3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_qc_lang1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k22_qc_lang1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_qc_lang1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_qc_lang1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0)) \Rightarrow (\forall X2.((\neg v1_xboole_0 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_qc_lang1 X0)))) \Rightarrow ((v5_qc_lang1 X1 X0) \Rightarrow (k4_qc_lang3 X0 X2 X1 = \\ k4_qc_lang3 X0 X2 (k22_qc_lang1 X0 X1)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0)) \Rightarrow (\forall X2.((\neg v1_xboole_0 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_qc_lang1 X0)))) \Rightarrow ((v3_qc_lang1 X1 X0) \Rightarrow (k4_qc_lang3 X0 X2 X1 = \\ k4_qc_lang3 X0 X2 (k18_qc_lang1 X0 X1)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0)) \Rightarrow ((v4_qc_lang2 X1 X0) \Rightarrow ((v3_qc_lang1 X1 X0) \wedge ((v5_qc_lang1 \\ (k18_qc_lang1 X0 X1) X0) \wedge (v3_qc_lang1 (k22_qc_lang1 X0 (k18_qc_lang1 \\ X0 X1)) X0)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((m1_qc_lang1 X0) \wedge (m1_subset_1 X1 (k9_qc_lang1 X0))) \Rightarrow (m1_subset_1 (k22_qc_lang1 X0 X1) (k9_qc_lang1 X0)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((m1_qc_lang1 X0) \wedge (m1_subset_1 X1 (k9_qc_lang1 X0))) \Rightarrow (m1_subset_1 (k18_qc_lang1 X0 X1) (k9_qc_lang1 X0)) \quad (5)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k9_qc_lang1 \\ X0)) \Rightarrow (\forall X2.((\neg v1_xboole_0\ X2) \wedge (m1_subset_1\ X2\ (k1_zfmisc_1 \\ (k2_qc_lang1\ X0)))) \Rightarrow ((v4_qc_lang2\ X1\ X0) \Rightarrow (k4_qc_lang3\ X0\ X2\ X1 = \\ k4_qc_lang3\ X0\ X2\ (k18_qc_lang1\ X0\ (k22_qc_lang1\ X0\ (k18_qc_lang1 \\ X0\ X1))))))) \end{aligned}$$