

t51_rinfsup1

(TMMe9VzFXXYXC3rb6ZcDX77RqJbYhebptoiW)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_seq_2 : \iota \Rightarrow o$ be given. Let $v8_valued_0 : \iota \Rightarrow o$ be given. Let $k4_rinfsup1 : \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k8_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m2_subset_1 X0 k1_numbers k5_numbers) \Rightarrow (\forall X1. \\ & ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers k1_numbers) \wedge (m1_subset_1 \\ & X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow ((v1_seq_2 \\ & X1) \Rightarrow (r1_xxreal_0 (k1_seq_1 (k4_rinfsup1 X1) (k2_nat_1 X0 np_1)) \\ & (k1_seq_1 (k4_rinfsup1 X1) X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(((v1_funct_1 X1) \wedge ((v1_funct_2 \\ & X1 k5_numbers X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\ & X0)))))) \wedge (v7_ordinal1 X2)) \Rightarrow (k8_nat_1 X0 X1 X2 = k1_funct_1 X1 X2) \end{aligned} \quad (4)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v3_valued_0 X0)))\Rightarrow(k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \quad (6)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (7)$$

Assume the following.

$$v3_membered k1_numbers \quad (8)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (9)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (10)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_funct_1 X0)\wedge((v1_funct_2 X0 k5_numbers k1_numbers)\wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers))))))\Rightarrow \\ & ((v1_funct_1 (k4_rinf sup1 X0))\wedge((v1_funct_2 (k4_rinf sup1 X0) \\ & k5_numbers k1_numbers)\wedge(m1_subset_1 (k4_rinf sup1 X0) (k1_zfmisc_1 \\ & (k2_zfmisc_1 k5_numbers k1_numbers)))))) \end{aligned} \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(m2_subset_1 (k2_nat_1 X0 X1) k1_numbers k5_numbers) \quad (12)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_funct_1 X0)\wedge((v1_funct_2 X0 k5_numbers k1_numbers)\wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers))))))\Rightarrow \\ & ((v8_valued_0 X0)\Leftrightarrow(\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers)\Rightarrow \\ & (r1_xxreal_0 (k8_nat_1 k1_numbers X0 (k2_nat_1 X1 np_1)) (k8_nat_1 \\ & k1_numbers X0 X1)))) \end{aligned} \quad (13)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.(v3_membered\ X1)\Rightarrow(\forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ X1)))\Rightarrow(v3_valued_0\ X2)) \quad (16)$$

Theorem 1

$$\forall X0.((v1_funct_1\ X0)\wedge((v1_funct_2\ X0\ k5_numbers\ k1_numbers)\wedge(m1_subset_1\ X0\ (k1_zfmisc_1\ (k2_zfmisc_1\ k5_numbers\ k1_numbers))))\Rightarrow((v1_seq_2\ X0)\Rightarrow(v8_valued_0\ (k4_rinf\ sup1\ X0)))$$