

t51_square_1 (TMS-
bXY25mbKVcds6yNkaUQX7zNGmDMTds5v)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_square_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (r1_xxreal_0 (k6_xcmplx_0 X0 X1) k6_numbers))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((r1_xxreal_0 (k6_xcmplx_0 (k3_square_1 X0) np_1) k6_numbers) \Rightarrow ((r1_xxreal_0 (k4_xcmplx_0 np_1) X0) \wedge (r1_xxreal_0 X0 np_1))) \quad (2)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (3)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (4)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xreal_0 (k3_square_1 X0)) \quad (5)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (6)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((r1_xxreal_0 (k3_square_1 X0) np_1) \Rightarrow ((r1_xxreal_0 (k4_xcmplx_0 np_1) X0) \wedge (r1_xxreal_0 X0 np_1)))$$