

t52\_afinsq\_1  
(TMWW3iw26eLoQf4Kzqip57dfTKMdMvGeVU6)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v5\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finset\_1 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_afinsq\_1 : \iota \Rightarrow \iota$  be given. Let  $k5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $r1\_ordinal1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k1\_card\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. (v1\_relat\_1 X1) \Rightarrow ((r1\_tarski (k9\_xtuple\_0 X1) X0) \Rightarrow (k5\_relat\_1 X1 X0 = X1)) \quad (1)$$

Assume the following.

$$\forall X0. (v7\_ordinal1 X0) \Rightarrow (\forall X1. (v7\_ordinal1 X1) \Rightarrow ((r1\_xxreal\_0 X0 X1) \Leftrightarrow (r1\_ordinal1 X0 X1))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v3\_ordinal1 X0) \wedge (v3\_ordinal1 X1)) \Rightarrow ((r1\_ordinal1 X0 X1) \Leftrightarrow (r1\_tarski X0 X1)) \quad (3)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v5\_ordinal1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finset\_1 X0)))) \Rightarrow (k1\_afinsq\_1 X0 = k1\_card\_1 X0) \quad (4)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v5\_ordinal1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finset\_1 X0)))) \Rightarrow (k1\_card\_1 X0 = k9\_xtuple\_0 X0) \quad (5)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v5\_ordinal1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finset\_1 X0)))) \Rightarrow (v7\_ordinal1 (k9\_xtuple\_0 X0)) \quad (6)$$

Assume the following.

$$\forall X0.(v5\_ordinal1 X0) \Leftrightarrow (v3\_ordinal1 (k9\_xtuple\_0 X0)) \quad (7)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (v3\_ordinal1 X0) \quad (8)$$

**Theorem 1**

$$\begin{aligned} \forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (( \\ v5\_ordinal1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finset\_1 X1)))) \Rightarrow ((r1\_xreal\_0 \\ (k1\_afinsq\_1 X1) X0) \Rightarrow (k5\_relat\_1 X1 X0 = X1))) \end{aligned}$$