

t52_euclid
(TMYtqutGcKZ4joDRSGxJ36j4yuEzoBQ2a6e)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k10_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $k7_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_finseq_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge ((v1_funct_1 X2) \wedge (v1_finseq_1 X2))) \Rightarrow ((X2 = k10_finseq_1 X0 X1) \Leftrightarrow ((k3_finseq_1 X2 = np_2) \wedge ((k1_funct_1 X2 np_1 = X0) \wedge (k1_funct_1 X2 np_2 = X1)))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v3_valued_0 X0))) \Rightarrow (k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (k19_euclid X0 X1 = k10_finseq_1 X0 X1) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 (k10_finseq_1 X0 X1)) \wedge (v1_funct_1 (k10_finseq_1 X0 X1)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (v3_valued_0 (k10_finseq_1 X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. v1_finseq_1 (k10_finseq_1 X0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(m1_subset_1 (k19_euclid X0 X1) (u1_struct_0 (k15_euclid np_2))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.k10_finseq_1 X0 X1 = k7_finseq_1 (k9_finseq_1 X0) (k9_finseq_1 X1) \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2)))\Rightarrow (k17_euclid X0 = k1_seq_1 X0 np_1) \quad (9)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2)))\Rightarrow (k18_euclid X0 = k1_seq_1 X0 np_2) \quad (10)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(\forall X1.(v1_xreal_0 X1)\Rightarrow((k17_euclid (k19_euclid X0 X1) = X0)\wedge(k18_euclid (k19_euclid X0 X1) = X1)))$$