

# t52\_fcont\_1 (TMVNubSXN- Qcx3BrZRda1mSfwobbCtpMinzN)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k1\_seq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_fcont\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xreal\_0 X2) \Rightarrow (\forall X3.(v1\_xreal\_0 X3) \Rightarrow (\neg(\neg r1\_xxreal\_0 \\ & X1 X0) \wedge ((r1\_xxreal\_0 X2 X3) \wedge (r1\_xxreal\_0 (k2\_xcmplx\_0 X1 X3) ( \\ & k2\_xcmplx\_0 X0 X2)))))))) \end{aligned} \quad (1)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xreal\_0 X2) \Rightarrow (\neg(\neg r1\_xxreal\_0 k6\_numbers X0) \wedge ((\neg r1\_xxreal\_0 \\ & X2 X1) \wedge (r1\_xxreal\_0 (k3\_xcmplx\_0 X1 X0) (k3\_xcmplx\_0 X2 X0)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0) \wedge (v1\_xreal\_0 X1)) \Rightarrow (v1\_xreal\_0 (k3\_xcmplx\_0 X0 X1)) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1\_xreal\_0 X0) \wedge (v1\_xreal\_0 X1)) \Rightarrow ((v1\_funct\_1 \\ & (k1\_fcont\_1 X0 X1)) \wedge ((v1\_funct\_2 (k1\_fcont\_1 X0 X1) k1\_numbers \\ & k1\_numbers) \wedge (m1\_subset\_1 (k1\_fcont\_1 X0 X1) (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & k1\_numbers k1\_numbers)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & ((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 X2 k1\_numbers k1\_numbers) \wedge (m1\_subset\_1 \\ & X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k1\_numbers k1\_numbers)))))) \Rightarrow ((X2 = \\ & k1\_fcont\_1 X0 X1) \Leftrightarrow (\forall X3.(v1\_xreal\_0 X3) \Rightarrow (k1\_seq\_1 X2 X3 = \\ & k2\_xcmplx\_0 (k3\_xcmplx\_0 X0 X3) X1)))))) \end{aligned} \quad (6)$$

Assume the following.

$$k1\_xboole\_0 = the (\lambda X0 : \iota.v1\_xboole\_0 X0) \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1\_xxreal\_0 X0) \wedge (v1\_xxreal\_0 X1)) \Rightarrow ( \\ & (r1\_xxreal\_0 X0 X1) \vee (r1\_xxreal\_0 X1 X0)) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1\_xcmplx\_0 X0) \wedge (v1\_xcmplx\_0 X1)) \Rightarrow ( \\ & k3\_xcmplx\_0 X0 X1 = k3\_xcmplx\_0 X1 X0) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xxreal\_0 X0) \quad (10)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xcmplx\_0 X0) \quad (11)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xreal\_0 X2) \Rightarrow (\forall X3.(v1\_xreal\_0 X3) \Rightarrow (\neg(\neg r1\_xxreal\_0 \\ & k6\_numbers X0) \wedge ((\neg r1\_xxreal\_0 X3 X2) \wedge (r1\_xxreal\_0 (k1\_seq\_1 \\ & (k1\_fcont\_1 X0 X1) X2) (k1\_seq\_1 (k1\_fcont\_1 X0 X1) X3))))))) \end{aligned}$$