

t52\_finseq\_2  
(TMZ1WKDWQNaU4BSTA5EGH9XM8mVfMKRLwAj)

October 27, 2020

Let  $k1\_finseq\_2 : \iota \Rightarrow \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $k10\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k1\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k12\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k5\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (k1\_finseq\_2 (k1\_nat\_1 X0 np\_1) = k7\_finseq\_1 (k1\_finseq\_2 X0) (k12\_finseq\_1 k5\_numbers (k1\_nat\_1 X0 np\_1))) \quad (1)$$

Assume the following.

$$k1\_finseq\_2 np\_1 = k12\_finseq\_1 k5\_numbers np\_1 \quad (2)$$

Assume the following.

$$((v2\_xxreal\_0 np\_2) \wedge (m2\_subset\_1 np\_2 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_2 k5\_numbers) \wedge (m1\_subset\_1 np\_2 k1\_numbers)) \quad (3)$$

Assume the following.

$$((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \quad (4)$$

Assume the following.

$$k2\_xcmplx\_0 np\_1 np\_1 = np\_2 \quad (5)$$

Assume the following.

$$\forall X0.k9\_finseq\_1 X0 = k5\_finseq\_1 X0 \quad (6)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v7\_ordinal1 X0)\wedge(m1\_subset\_1 X1 k5\_numbers))\Rightarrow (k1\_nat\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0)\wedge(m1\_subset\_1 X1 X0))\Rightarrow (k12\_finseq\_1 X0 X1 = k5\_finseq\_1 X1) \quad (9)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1)\wedge(v3\_ordinal1 k4\_ordinal1) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.k10\_finseq\_1 X0 X1 = k7\_finseq\_1 (k9\_finseq\_1 X0) (k9\_finseq\_1 X1) \quad (11)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1)\Rightarrow(v7\_ordinal1 X0) \quad (12)$$

**Theorem 1**  $k1\_finseq\_2 np\_2 = k10\_finseq\_1 np\_1 np\_2$ .