

t52_glib_003

(TMTK7qnQNKtCmxRTMU1TYns4bfE6i3JMiCd)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_glib_000 : \iota \Rightarrow o$ be given. Let $v3_glib_003 : \iota \Rightarrow o$ be given. Let $k6_glib_000 : \iota \Rightarrow \iota$ be given. Let $k14_glib_003 : \iota \Rightarrow \iota$ be given. Let $k13_glib_003 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k7_glib_003 : \iota \Rightarrow \iota$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k7_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge ((v1_glib_000 X0) \wedge (v3_glib_003 X0)))))) \Rightarrow \\ (\forall X1. \forall X2. (X1 \in k6_glib_000 X0) \Rightarrow (k7_glib_003 (k13_glib_003 \\ X0 X1 X2) = k1_funct_4 (k7_glib_003 X0) (k16_funcop_1 X1 X2))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (k9_xtuple_0 (k2_funcop_1 X0 X1) = X0) \wedge (r1_tarski (k10_xtuple_0 (k2_funcop_1 X0 X1)) (k1_tarski X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. k7_funcop_1 X0 X1 = k2_funcop_1 X0 X1 \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. v1_relat_1 (k2_zfmisc_1 X0 X1) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge ((v1_glib_000 X0) \wedge (v3_glib_003 X0)))))) \Rightarrow \\ ((v1_relat_1 (k7_glib_003 X0)) \wedge (v1_funct_1 (k7_glib_003 X0))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(v1_funct_1 (k7_funcop_1 X0 X1))\wedge((v1_funct_2 (k7_funcop_1 X0 X1) X0 (k1_tarSKI X1))\wedge(m1_subset_1 (k7_funcop_1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 (k1_tarSKI X1)))))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(((v1_relat_1 X0)\wedge(v1_funct_1 X0))\wedge((v1_relat_1 X1)\wedge(v1_funct_1 X1)))\Rightarrow((v1_relat_1 (k1_funct_4 X0 X1))\wedge(v1_funct_1 (k1_funct_4 X0 X1))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1_relat_1 X0)\wedge((v4_relat_1 X0 k5_numbers)\wedge((v1_funct_1 X0)\wedge((v1_finset_1 X0)\wedge((v1_glib_000 X0)\wedge(v3_glib_003 X0))))))\Rightarrow((v1_relat_1 (k13_glib_003 X0 X1 X2))\wedge((v4_relat_1 (k13_glib_003 X0 X1 X2) k5_numbers)\wedge((v1_funct_1 (k13_glib_003 X0 X1 X2))\wedge((v1_finset_1 (k13_glib_003 X0 X1 X2))\wedge((v1_glib_000 (k13_glib_003 X0 X1 X2))\wedge(v3_glib_003 (k13_glib_003 X0 X1 X2)))))))) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.k16_funcop_1 X0 X1 = k7_funcop_1 (k1_tarSKI X0) X1 \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.k2_funcop_1 X0 X1 = k2_zfmisc_1 X0 (k1_tarSKI X1) \quad (10)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v4_relat_1 X0 k5_numbers)\wedge((v1_funct_1 X0)\wedge((v1_finset_1 X0)\wedge((v1_glib_000 X0)\wedge(v3_glib_003 X0))))))\Rightarrow (k14_glib_003 X0 = k9_xtuple_0 (k7_glib_003 X0)) \quad (11)$$

Assume the following.

$$\forall X0.(((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(\forall X1.(((v1_relat_1 X1)\wedge(v1_funct_1 X1))\Rightarrow(\forall X2.(((v1_relat_1 X2)\wedge(v1_funct_1 X2))\Rightarrow((X2 = k1_funct_4 X0 X1)\Leftrightarrow((k9_xtuple_0 X2 = k2_xboole_0 (k9_xtuple_0 X0) (k9_xtuple_0 X1))\wedge(\forall X3.(X3 \in k2_xboole_0 (k9_xtuple_0 X0) (k9_xtuple_0 X1))\Rightarrow(((X3 \in k9_xtuple_0 X1)\Rightarrow(k1_funct_1 X2 X3 = k1_funct_1 X1 X3))\wedge((\neg X3 \in k9_xtuple_0 X1)\Rightarrow(k1_funct_1 X2 X3 = k1_funct_1 X0 X3)))))))))) \quad (12)$$

Theorem 1

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 \ k5_numbers) \wedge ((v1_funct_1 \\ & X0) \wedge ((v1_finset_1 X0) \wedge ((v1_glib_000 X0) \wedge (v3_glib_003 X0)))))) \Rightarrow \\ & (\forall X1. \forall X2. (X1 \in k6_glib_000 X0) \Rightarrow (k14_glib_003 (k13_glib_003 \\ & X0 X1 X2) = k2_xboole_0 (k14_glib_003 X0) (k1_tarski X1))) \end{aligned}$$