

t52_nat_1

(TMYXpn1edH94gbWugfZtgda1crtwHjqtrcc)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k8_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \wedge (v7_ordinal1 X2)) \Rightarrow (k8_nat_1 X0 X1 X2 = k1_funct_1 X1 X2) \quad (1)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (3)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. (X1 = k10_xtuple_0 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. (X3 \in k9_xtuple_0 X0) \wedge (X2 = k1_funct_1 X0 X3)))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow ((v4_relat_1 X1 X0) \Leftrightarrow (r1_tarski (k9_xtuple_0 X1) X0)) \quad (5)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Leftrightarrow (X0 \in k4_ordinal1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (8)$$

Theorem 1

$$\forall X0.\forall X1.(\neg v1_xboole_0 X1)\Rightarrow(\forall X2.((v1_funct_1 X2)\wedge((v1_funct_2 X2 k5_numbers X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X1)))))\Rightarrow((\forall X3.(v7_ordinal1 X3)\Rightarrow(k8_nat_1 X1 X2 X3 \in X0))\Rightarrow(r1_tarSKI (k10_xtuple_0 X2) X0)))$$