

t52\_xcplx\_1  
(TMNLT9BhcoiRjLiEEqZxxeBF5VZCcJz42v9)

October 27, 2020

Let  $v1\_xcplx\_0 : \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k7\_xcplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_xcplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1\_xcplx\_0 X0) \wedge ((v1\_xcplx\_0 X1) \wedge (v1\_xcplx\_0 X2))) \Rightarrow (k3\_xcplx\_0 X0 (k7\_xcplx\_0 X1 X2) = k7\_xcplx\_0 (k3\_xcplx\_0 X0 X1) X2) \quad (1)$$

Assume the following.

$$\forall X0. (v1\_xcplx\_0 X0) \Rightarrow (\forall X1. (v1\_xcplx\_0 X1) \Rightarrow ((X0 \neq k6\_numbers) \Rightarrow (k3\_xcplx\_0 (k7\_xcplx\_0 X1 X0) X0 = X1))) \quad (2)$$

Assume the following.

$$\forall X0. (v1\_xcplx\_0 X0) \Rightarrow (\forall X1. (v1\_xcplx\_0 X1) \Rightarrow (\forall X2. (v1\_xcplx\_0 X2) \Rightarrow (k7\_xcplx\_0 X0 (k7\_xcplx\_0 X1 X2) = k7\_xcplx\_0 (k3\_xcplx\_0 X0 X2) X1))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_xcplx\_0 X0) \wedge (v1\_xcplx\_0 X1)) \Rightarrow (v1\_xcplx\_0 (k7\_xcplx\_0 X0 X1)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_xcplx\_0 X0) \wedge (v1\_xcplx\_0 X1)) \Rightarrow (k3\_xcplx\_0 X0 X1 = k3\_xcplx\_0 X1 X0) \quad (5)$$

**Theorem 1**

$$\forall X0. (v1\_xcplx\_0 X0) \Rightarrow (\forall X1. (v1\_xcplx\_0 X1) \Rightarrow ((X0 \neq k6\_numbers) \Rightarrow (k7\_xcplx\_0 X0 (k7\_xcplx\_0 X0 X1) = X1)))$$