

t53_card_2 (TMFKbobDJwRVFt- GknPbHJ9BMcDyb62LFoJt)

October 27, 2020

Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_card_1 : \iota \Rightarrow \iota$ be given. Let $k3_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_5 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k2_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_4 : \iota$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (\forall X3.(v1_xreal_0 X3) \Rightarrow (((r1_xxreal_0 \\ & X0 X1) \wedge (r1_xxreal_0 X2 X3)) \Rightarrow (r1_xxreal_0 (k2_xcmplx_0 X0 X2) (\\ & k2_xcmplx_0 X1 X3)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.\forall X4.k3_enumset1 X0 X1 X2 X3 X4 = k2_xboole_0 (k1_tarski X0) (k2_enumset1 X1 X2 X3 X4) \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.r1_xxreal_0 (k5_card_1 (k2_enumset1 X0 X1 X2 X3)) np_4 \tag{3}$$

Assume the following.

$$\forall X0.(v1_finset_1 X0) \Rightarrow (\forall X1.(v1_finset_1 X1) \Rightarrow (r1_xreal_0 (k5_card_1 (k2_xboole_0 X0 X1)) (k2_nat_1 (k5_card_1 X0) (k5_card_1 X1)))) \tag{4}$$

Assume the following.

$$\forall X0.k1_card_1 (k1_tarski X0) = np_1 \tag{5}$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xreal_0 X0 X1) \wedge (r1_xreal_0 X1 X2)) \Rightarrow \\ & (r1_xreal_0 X0 X2)))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & ((v2_xreal_0 np_5) \wedge (m2_subset_1 np_5 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_5 k5_numbers) \wedge (m1_subset_1 np_5 k1_numbers)) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & ((v2_xreal_0 np_4) \wedge (m2_subset_1 np_4 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_4 k5_numbers) \wedge (m1_subset_1 np_4 k1_numbers)) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & ((v2_xreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (9)$$

Assume the following.

$$k2_xcmplx_0 np_1 np_4 = np_5 \quad (10)$$

Assume the following.

$$r1_xreal_0 np_1 np_1 \quad (11)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (12)$$

Assume the following.

$$\forall X0.(v1_finset_1 X0) \Rightarrow (k5_card_1 X0 = k1_card_1 X0) \quad (13)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((m1_subset_1 X0 k5_numbers) \wedge (v7_ordinal1 \\ & X1)) \Rightarrow (k2_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \end{aligned} \quad (14)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (v1_xreal_0 \\ & (k2_xcmplx_0 X0 X1)) \end{aligned} \quad (15)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.\forall X4.v1_finset_1 \\ & (k3_enumset1 X0 X1 X2 X3 X4) \end{aligned} \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.v1_finset_1 (k2_enumset1 X0 X1 X2 X3) \quad (17)$$

Assume the following.

$$\forall X0.v1_finset_1 (k1_tarski X0) \quad (18)$$

Assume the following.

$$\forall X0.(v1_finset_1 X0)\Rightarrow(m1_subset_1 (k5_card_1 X0) k4_ordinal1) \quad (19)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (20)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xxreal_0 X0) \quad (21)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(v1_xreal_0 X0) \quad (22)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (23)$$

Theorem 1

$$\forall X0.\forall X1.\forall X2.\forall X3.\forall X4.r1_xxreal_0 (k5_card_1 (k3_enumset1 X0 X1 X2 X3 X4)) np_5$$