

t53_seq_1

(TMWZRk6V6CoXr9SNQjRcF7Yh2htX7A3vaWE)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_relat_1 : \iota \Rightarrow o$ be given. Let $k56_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k17_complex1 : \iota \Rightarrow \iota$ be given. Let $r2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k18_complex1 : \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_membered : \iota \Rightarrow o$ be given. Let $k54_valued_1 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k16_complex1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. ((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\ & ((v2_relat_1 X0) \Leftrightarrow (\forall X1. (m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow \\ & (k1_seq_1 X0 X1 \neq k6_numbers))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. (v1_xcmplx_0 X0) \Rightarrow ((\neg(X0 \neq k6_numbers) \wedge (r1_xxreal_0 \\ & (k17_complex1 X0) k6_numbers)) \wedge (\neg(\neg r1_xxreal_0 (k17_complex1 \\ & X0) k6_numbers) \wedge (X0 = k6_numbers))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned}
& \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\
& (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\
& (\forall X1.((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers k1_numbers) \wedge \\
& (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\
& ((r2_funct_2 k5_numbers k1_numbers X0 (k56_valued_1 k5_numbers \\
& k1_numbers X1)) \Leftrightarrow (\forall X2.(m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow \\
& (k1_seq_1 X0 X2 = k18_complex1 (k1_seq_1 X1 X2))))))
\end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned}
& \forall X0. \forall X1. \forall X2. \forall X3. (((v1_funct_1 X2) \wedge \\
& ((v1_funct_2 X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\
& X0 X1)))))) \wedge ((v1_funct_1 X3) \wedge ((v1_funct_2 X3 X0 X1) \wedge (m1_subset_1 \\
& X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \Rightarrow (r2_funct_2 X0 X1 X2 X2)
\end{aligned} \tag{4}$$

Assume the following.

$$k6_numbers = k1_xboole_0 \tag{5}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{6}$$

Assume the following.

$$\begin{aligned}
& \forall X0. \forall X1. \forall X2. ((v1_membered X1) \wedge ((v1_funct_1 \\
& X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \Rightarrow (k56_valued_1 \\
& X0 X1 X2 = k54_valued_1 X2)
\end{aligned} \tag{7}$$

Assume the following.

$$\begin{aligned}
& \forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v3_valued_0 \\
& X0))) \Rightarrow (k1_seq_1 X0 X1 = k1_funct_1 X0 X1)
\end{aligned} \tag{8}$$

Assume the following.

$$\forall X0. (v1_xcmplx_0 X0) \Rightarrow (k18_complex1 X0 = k16_complex1 X0) \tag{9}$$

Assume the following.

$$\forall X0. (v1_xcmplx_0 X0) \Rightarrow (k17_complex1 X0 = k16_complex1 X0) \tag{10}$$

Assume the following.

$$\begin{aligned}
& \forall X0. (v1_xcmplx_0 X0) \Rightarrow (k17_complex1 (k17_complex1 X0) = \\
& k17_complex1 X0)
\end{aligned} \tag{11}$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(((\neg v1_xboole_0 X1)\wedge(v1_membered \\ X1))\wedge((v1_funct_1 X2)\wedge((v1_funct_2 X2 X0 X1)\wedge(m1_subset_1 X2 \\ (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))))\Rightarrow((v1_funct_1 (k54_valued_1 \\ X2))\wedge(v1_partfun1 (k54_valued_1 X2) X0)) \end{aligned} \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_valued_0 \\ X0)))\Rightarrow(v1_xcmplx_0 (k1_funct_1 X0 X1)) \quad (13)$$

Assume the following.

$$v3_membered k1_numbers \quad (14)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (15)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((v1_membered X1)\wedge((v1_funct_1 \\ X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))\Rightarrow((v1_funct_1 \\ (k56_valued_1 X0 X1 X2))\wedge(m1_subset_1 (k56_valued_1 X0 X1 X2) (\\ k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \end{aligned} \quad (16)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0)\Rightarrow(v1_xreal_0 (k16_complex1 X0)) \quad (17)$$

Assume the following.

$$\forall X0.(v3_membered X0)\Rightarrow(v1_membered X0) \quad (18)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xcmplx_0 X0) \quad (19)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (20)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1)))\Rightarrow((v1_partfun1 X2 X0)\Rightarrow(v1_funct_2 X2 X0 X1)) \end{aligned} \quad (21)$$

Assume the following.

$$\forall X0.\forall X1.(v3_membered X1)\Rightarrow(\forall X2.(m1_subset_1 \\ X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v3_valued_0 X2)) \quad (22)$$

Assume the following.

$$\forall X0.\forall X1.(v1_membered\ X1)\Rightarrow(\forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ X1)))\Rightarrow(v1_valued_0\ X2)) \quad (23)$$

Theorem 1

$$\forall X0.((v1_funct_1\ X0)\wedge((v1_funct_2\ X0\ k5_numbers\ k1_numbers)\wedge(m1_subset_1\ X0\ (k1_zfmisc_1\ (k2_zfmisc_1\ k5_numbers\ k1_numbers))))\Rightarrow((v2_relat_1\ X0)\Rightarrow(v2_relat_1\ (k56_valued_1\ k5_numbers\ k1_numbers\ X0)))$$