

t53_setlim_1 (TMZyUzxAduRJAfT-
wau7zu2xk22GTcWqmAMx)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k9_setfam_1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_prob_1 : \iota \Rightarrow o$ be given. Let $k1_kurato_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_setlim_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_prob_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k5_numbers) \Rightarrow (\forall X1.\forall X2. \\ & ((v1_funct_1 X2) \wedge ((v1_funct_2 X2 k5_numbers (k9_setfam_1 X1)) \wedge \\ & (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (k9_setfam_1 \\ & X1)))))) \Rightarrow ((v2_prob_1 X2) \Rightarrow (k3_funct_2 k5_numbers (k9_setfam_1 \\ & X1) (k2_setlim_1 X1 X2) X0 = k3_prob_1 X1 X2))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (\forall X2. \\ & ((v1_funct_1 X2) \wedge ((v1_funct_2 X2 k5_numbers (k9_setfam_1 X0)) \wedge \\ & (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (k9_setfam_1 \\ & X0)))))) \Rightarrow ((\forall X3.(m1_subset_1 X3 k5_numbers) \Rightarrow (k3_funct_2 \\ & k5_numbers (k9_setfam_1 X0) X2 X3 = X1)) \Rightarrow (k1_kurato_0 X0 X2 = X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.k9_setfam_1 X0 = k1_zfmisc_1 X0 \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers \\ & (k9_setfam_1 X0)) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers (k9_setfam_1 X0)))))) \Rightarrow (m1_subset_1 (k3_prob_1 X0 \\ & X1) (k1_zfmisc_1 X0)) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers \\ & (k9_setfam_1 X0)) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers (k9_setfam_1 X0)))))) \Rightarrow ((v1_funct_1 (k2_setlim_1 \\ & X0 X1)) \wedge ((v1_funct_2 (k2_setlim_1 X0 X1) k5_numbers (k9_setfam_1 \\ & X0)) \wedge (m1_subset_1 (k2_setlim_1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers (k9_setfam_1 X0)))))) \end{aligned} \tag{5}$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers \\ & (k9_setfam_1 X0)) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers (k9_setfam_1 X0)))))) \Rightarrow ((v2_prob_1 X1) \Rightarrow (k1_kurato_0 \\ & X0 (k2_setlim_1 X0 X1) = k3_prob_1 X0 X1)) \end{aligned}$$