

t53_zfmisc_1

(TMaCZ64gM97kePruziRFgCHgxoH4mzE1TKv)

October 27, 2020

Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k3_xboole_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (X3 \in X1))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k2_tarski X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 = X0) \vee (X3 = X1))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (3)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. \neg (k3_xboole_0 (k2_tarski X0 X1) X2 = k1_tarski X0) \wedge ((X1 \in X2) \wedge (X0 \neq X1))$$