

t54_rfunct_1

(TMYLKVKAZEKK9HW9wn9pBcbDSr4PLTQmNj3)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_rfunct_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $k4_rfunct_1 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarski X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. k4_xboole_0 X0 k1_xboole_0 = X0 \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 (k4_xboole_0 X1 X0)) \Rightarrow (X0 = k1_xboole_0) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (k8_relset_1 X0 X1 X2 X3 = k8_relat_1 X2 X3) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. k6_subset_1 X0 X1 = k4_xboole_0 X0 X1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v3_membered\ X1)\wedge((v1_funct_1\ X2)\wedge(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ X1))))))\Rightarrow(k6_rfunct_1\ X0\ X1\ X2 = k4_rfunct_1\ X2) \quad (6)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1\ X1)\wedge(v4_relat_1\ X1\ X0))\Rightarrow(k1_relset_1\ X0\ X1 = k9_xtuple_0\ X1) \quad (8)$$

Assume the following.

$$v3_membered\ k1_numbers \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ X1)))\Rightarrow(m1_subset_1\ (k8_relset_1\ X0\ X1\ X2\ X3)\ (k1_zfmisc_1\ X0)) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v3_membered\ X1)\wedge((v1_funct_1\ X2)\wedge(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ X1))))))\Rightarrow((v1_funct_1\ (k6_rfunct_1\ X0\ X1\ X2))\wedge(m1_subset_1\ (k6_rfunct_1\ X0\ X1\ X2)\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ k1_numbers)))) \quad (11)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0)\wedge((v1_funct_1\ X0)\wedge(v1_valued_0\ X0)))\Rightarrow(\forall X1.((v1_relat_1\ X1)\wedge(v1_funct_1\ X1))\Rightarrow((X1 = k4_rfunct_1\ X0)\Leftrightarrow((k9_xtuple_0\ X1 = k6_subset_1\ (k9_xtuple_0\ X0)\ (k8_relat_1\ X0\ (k1_tarski\ k6_numbers))))\wedge(\forall X2.(X2 \in k9_xtuple_0\ X1)\Rightarrow(k1_funct_1\ X1\ X2 = k5_xcmplx_0\ (k1_funct_1\ X0\ X2)))))) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1\ X1)\wedge(v4_relat_1\ X1\ X0))\Rightarrow((v1_partfun1\ X1\ X0)\Leftrightarrow(k1_relset_1\ X0\ X1 = X0)) \quad (13)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0)\wedge(v3_valued_0\ X0))\Rightarrow((v1_relat_1\ X0)\wedge(v1_valued_0\ X0)) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.(v3_membered X1)\Rightarrow(\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v3_valued_0 X2)) \quad (17)$$

Theorem 1

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((v1_funct_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers))))\Rightarrow((v1_partfun1 (k6_rfunct_1 X0 k1_numbers X1) X0)\Leftrightarrow((k8_relset_1 X0 k1_numbers X1 (k1_tarski k6_numbers) = k1_xboole_0)\wedge(v1_partfun1 X1 X0))))$$