

t54_sin_cos

(TMGH5nkH12WytijRCLmuYoXwnxFh1xPL8Jw)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k24_sin_cos : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee ((v3_xxreal_0 X0) \vee (v2_xxreal_0 X1)))))) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((\neg r1_xxreal_0 k6_numbers X0) \Rightarrow ((\neg r1_xxreal_0 (k1_seq_1 k24_sin_cos X0) k6_numbers) \wedge (r1_xxreal_0 (k1_seq_1 k24_sin_cos X0) np_1))) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((\neg r1_xxreal_0 X0 k6_numbers) \Rightarrow (r1_xxreal_0 np_1 (k1_seq_1 k24_sin_cos X0))) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (v2_xxreal_0 X0)) \Rightarrow (v2_xxreal_0 X1))) \quad (5)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 \ np_1) \wedge (m2_subset_1 \ np_1 \ k1_numbers \ k5_numbers)) \wedge \\ & ((m1_subset_1 \ np_1 \ k5_numbers) \wedge (m1_subset_1 \ np_1 \ k1_numbers)) \end{aligned} \quad (6)$$

Assume the following.

$$v1_xboole_0 \ np_0 \quad (7)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (8)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 \ X0) \wedge ((v1_funct_1 \ X0) \wedge (v3_valued_0 \ X0))) \Rightarrow (k1_seq_1 \ X0 \ X1 = k1_funct_1 \ X0 \ X1) \quad (9)$$

Assume the following.

$$\exists X0. (v1_xboole_0 \ X0) \wedge ((v1_xcmplx_0 \ X0) \wedge ((v1_xxreal_0 \ X0) \wedge (v1_xxreal_0 \ X0))) \quad (10)$$

Assume the following.

$$k1_seq_1 \ k24_sin_cos \ k6_numbers = np_1 \quad (11)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 \ X0) \wedge ((v1_funct_1 \ X0) \wedge (v3_valued_0 \ X0))) \Rightarrow (v1_xreal_0 \ (k1_funct_1 \ X0 \ X1)) \quad (12)$$

Assume the following.

$$\begin{aligned} & (v1_funct_1 \ k24_sin_cos) \wedge ((v1_funct_2 \ k24_sin_cos \ k1_numbers \\ & k1_numbers) \wedge (m1_subset_1 \ k24_sin_cos \ (k1_zfmisc_1 \ (k2_zfmisc_1 \\ & k1_numbers \ k1_numbers)))) \end{aligned} \quad (13)$$

Assume the following.

$$\forall X0. ((v1_xxreal_0 \ X0) \wedge (v3_xxreal_0 \ X0)) \Rightarrow ((\neg v1_xboole_0 \ X0) \wedge ((v1_xxreal_0 \ X0) \wedge (\neg v2_xxreal_0 \ X0))) \quad (14)$$

Assume the following.

$$\forall X0. (v1_xreal_0 \ X0) \Rightarrow (v1_xxreal_0 \ X0) \quad (15)$$

Assume the following.

$$\forall X0. ((v1_xxreal_0 \ X0) \wedge (v2_xxreal_0 \ X0)) \Rightarrow ((\neg v1_xboole_0 \ X0) \wedge ((v1_xxreal_0 \ X0) \wedge (\neg v3_xxreal_0 \ X0))) \quad (16)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v5_relat_1 X0 k1_numbers)) \Rightarrow ((v1_relat_1 X0) \wedge (v3_valued_0 X0)) \quad (17)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \quad (18)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (19)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (20)$$

Theorem 1

$$\forall X0. (v1_xreal_0 X0) \Rightarrow (\neg r1_xxreal_0 (k1_seq_1 k24_sin_cos X0) k6_numbers)$$