

t54_xreal_1

(TMG2Deanr1vML1Lg9PLxrEFbcC9m8ZpH8vG)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 k6_numbers X0) \wedge (r1_xxreal_0 (k6_xcmplx_0 X1 X0) X1))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2.(v1_xreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow (r1_xxreal_0 X0 X2)))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (v1_xreal_0 (k6_xcmplx_0 X0 X1)) \quad (3)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2.(v1_xreal_0 X2) \Rightarrow (\neg(\neg r1_xxreal_0 k6_numbers X0) \wedge ((r1_xxreal_0 X1 X2) \wedge (r1_xxreal_0 (k6_xcmplx_0 X2 X0) X1))))))$$