

t55_afinsq_1
(TMHjWtfH8o4t5oeHivfGq8smNnPYd1hbbTb)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v5_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_afinsq_1 : \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge ((\\ v5_ordinal1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finset_1 X1)))) \Rightarrow ((r1_xxreal_0 \\ X0 (k1_afinsq_1 X1)) \Rightarrow (k1_afinsq_1 (k5_relat_1 X1 X0) = X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge ((\\ v5_ordinal1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finset_1 X1)))) \Rightarrow ((r1_xxreal_0 \\ (k1_afinsq_1 X1) X0) \Rightarrow (k5_relat_1 X1 X0 = X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (\\ r1_xxreal_0 X0 X0) \quad (3)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v5_ordinal1 X0) \wedge ((v1_funct_1 \\ X0) \wedge (v1_finset_1 X0)))) \Rightarrow (k1_afinsq_1 X0 = k1_card_1 X0) \quad (4)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v5_ordinal1 X0) \wedge ((v1_funct_1 \\ X0) \wedge (v1_finset_1 X0)))) \Rightarrow (k1_card_1 X0 = k9_xtuple_0 X0) \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v5_ordinal1 X0) \wedge ((v1_funct_1 \\ X0) \wedge (v1_finset_1 X0)))) \Rightarrow (v7_ordinal1 (k9_xtuple_0 X0)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0)\wedge(v1_xxreal_0 X1))\Rightarrow((r1_xxreal_0 X0 X1)\vee(r1_xxreal_0 X1 X0)) \quad (7)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(v1_xxreal_0 X0) \quad (8)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(\forall X1.((v1_relat_1 X1)\wedge((v5_ordinal1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finset_1 X1))))\Rightarrow(r1_xxreal_0 (k1_afinsq_1 (k5_relat_1 X1 X0)) X0))$$