

t55_compos_2 (TMQP- TjLC3z1LZA3jsuPmktbwBwRRkcQSQb)

October 27, 2020

Let $v1_amistd_4 : \iota \Rightarrow o$ be given. Let $l1_compos_1 : \iota \Rightarrow o$ be given. Let $v6_compos_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_compos_1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_compos_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_compos_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_3 : \iota$ be given. Let $k5_compos_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_afinsq_1 : \iota \Rightarrow o$ be given. Let $v3_compos_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_compos_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k62_valued_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned}
& \forall X0.((v1_amistd_4 X0) \wedge (l1_compos_1 X0)) \Rightarrow (\forall X1. \\
& ((v6_compos_0 X1 (u1_compos_1 X0)) \wedge (m1_subset_1 X1 (u1_compos_1 \\
& X0))) \Rightarrow (\forall X2.((\neg v1_xboole_0 X2) \wedge ((v1_relat_1 X2) \wedge ((v4_relat_1 \\
& X2 k5_numbers) \wedge ((v5_relat_1 X2 (u1_compos_1 X0)) \wedge ((v1_funct_1 \\
& X2) \wedge ((v1_finset_1 X2) \wedge ((v1_afinsq_1 X2) \wedge ((v3_compos_1 X2 X0) \wedge \\
& (v4_compos_1 X2 X0)))))))))) \Rightarrow (k1_funct_1 (k2_compos_2 X0 X2 X1) \\
& (k62_valued_1 X2) = k5_compos_0 (u1_compos_1 X0) X1 (k62_valued_1 \\
& X2)))
\end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
& \forall X0.((v1_amistd_4 X0) \wedge (l1_compos_1 X0)) \Rightarrow (\forall X1. \\
& ((v6_compos_0 X1 (u1_compos_1 X0)) \wedge (m1_subset_1 X1 (u1_compos_1 \\
& X0))) \Rightarrow (\forall X2.((v6_compos_0 X2 (u1_compos_1 X0)) \wedge (m1_subset_1 \\
& X2 (u1_compos_1 X0))) \Rightarrow (\forall X3.((v6_compos_0 X3 (u1_compos_1 \\
& X0)) \wedge (m1_subset_1 X3 (u1_compos_1 X0))) \Rightarrow (k62_valued_1 (k2_compos_2 \\
& X0 (k3_compos_2 X0 X1 X2) X3) = np_3)))
\end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned}
& \forall X0.\forall X1.\forall X2.(((v1_amistd_4 X0)\wedge(l1_compos_1 \\
& X0))\wedge(((v6_compos_0 X1 (u1_compos_1 X0))\wedge(m1_subset_1 X1 (u1_compos_1 \\
& X0))\wedge((v6_compos_0 X2 (u1_compos_1 X0))\wedge(m1_subset_1 X2 (u1_compos_1 \\
& X0))))))\Rightarrow((\neg v1_xboole_0 (k3_compos_2 X0 X1 X2))\wedge((v1_relat_1 \\
& (k3_compos_2 X0 X1 X2))\wedge((v4_relat_1 (k3_compos_2 X0 X1 X2) k5_numbers)\wedge \\
& ((v5_relat_1 (k3_compos_2 X0 X1 X2) (u1_compos_1 X0))\wedge((v1_funct_1 \\
& (k3_compos_2 X0 X1 X2))\wedge((v1_finset_1 (k3_compos_2 X0 X1 X2))\wedge \\
& ((v1_afinsq_1 (k3_compos_2 X0 X1 X2))\wedge((v3_compos_1 (k3_compos_2 \\
& X0 X1 X2) X0)\wedge(v4_compos_1 (k3_compos_2 X0 X1 X2) X0)))))))))) \\
& \tag{3}
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& \forall X0.\forall X1.\forall X2.(((v1_amistd_4 X0)\wedge(l1_compos_1 \\
& X0))\wedge(((\neg v1_xboole_0 X1)\wedge((v1_relat_1 X1)\wedge((v4_relat_1 X1 k5_numbers)\wedge \\
& ((v5_relat_1 X1 (u1_compos_1 X0))\wedge((v1_funct_1 X1)\wedge((v1_finset_1 \\
& X1)\wedge((v1_afinsq_1 X1)\wedge((v3_compos_1 X1 X0)\wedge(v4_compos_1 X1 X0))))))))))\wedge \\
& ((v6_compos_0 X2 (u1_compos_1 X0))\wedge(m1_subset_1 X2 (u1_compos_1 \\
& X0))))))\Rightarrow((\neg v1_xboole_0 (k2_compos_2 X0 X1 X2))\wedge((v1_relat_1 \\
& (k2_compos_2 X0 X1 X2))\wedge((v4_relat_1 (k2_compos_2 X0 X1 X2) k5_numbers)\wedge \\
& ((v5_relat_1 (k2_compos_2 X0 X1 X2) (u1_compos_1 X0))\wedge((v1_funct_1 \\
& (k2_compos_2 X0 X1 X2))\wedge((v1_finset_1 (k2_compos_2 X0 X1 X2))\wedge \\
& ((v1_afinsq_1 (k2_compos_2 X0 X1 X2))\wedge((v3_compos_1 (k2_compos_2 \\
& X0 X1 X2) X0)\wedge(v4_compos_1 (k2_compos_2 X0 X1 X2) X0)))))))))) \\
& \tag{4}
\end{aligned}$$

Theorem 1

$$\begin{aligned}
& \forall X0.((v1_amistd_4 X0)\wedge(l1_compos_1 X0))\Rightarrow(\forall X1. \\
& ((v6_compos_0 X1 (u1_compos_1 X0))\wedge(m1_subset_1 X1 (u1_compos_1 \\
& X0)))\Rightarrow(\forall X2.((v6_compos_0 X2 (u1_compos_1 X0))\wedge(m1_subset_1 \\
& X2 (u1_compos_1 X0)))\Rightarrow(\forall X3.((v6_compos_0 X3 (u1_compos_1 \\
& X0))\wedge(m1_subset_1 X3 (u1_compos_1 X0)))\Rightarrow(\forall X4.((v6_compos_0 \\
& X4 (u1_compos_1 X0))\wedge(m1_subset_1 X4 (u1_compos_1 X0)))\Rightarrow(k1_funct_1 \\
& (k2_compos_2 X0 (k2_compos_2 X0 (k3_compos_2 X0 X1 X2) X3) X4) np_3 = \\
& k5_compos_0 (u1_compos_1 X0) X4 np_3))))))
\end{aligned}$$