

t55_flang_3 (TMSBswe-
QbGNpQ533oLzUQ3RSY55Nji7hShk)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k8_afinsq_1 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_flang_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_flang_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_catalan2 : \iota \Rightarrow \iota$ be given. Let $k6_flang_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k3_catalan2 X0))) \Rightarrow ((r1_tarski (k6_flang_1 X0 (k8_flang_1 X0 X1) X1) (k8_flang_1 X0 X1)) \wedge (r1_tarski (k6_flang_1 X0 X1 (k8_flang_1 X0 X1)) (k8_flang_1 X0 X1))) \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 X0))) \Rightarrow (k2_flang_3 X0 X1 = k6_flang_1 X0 (k8_flang_1 X0 X1) X1) \tag{2}$$

Assume the following.

$$\forall X0. k3_catalan2 X0 = k8_afinsq_1 X0 \tag{3}$$

Theorem 1

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k8_afinsq_1 X0))) \Rightarrow (r1_tarski (k2_flang_3 X0 X1) (k8_flang_1 X0 X1))$$