

t55_partfun1
(TMdC3MsAG1BEiMBXLKoXb8VKquhHP2k8vS3)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $r1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_partfun1 X0 X1) \Leftrightarrow (\exists X2. \\ ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \wedge ((r1_tarski X0 X2) \wedge (r1_tarski \\ X1 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_tarski X0 X1) \Rightarrow (k5_relat_1 \\ X1 (k9_xtuple_0 X0) = X0))) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (((k9_xtuple_0 X0 = k9_xtuple_0 \\ X1) \wedge (r1_partfun1 X0 X1)) \Rightarrow (X0 = X1))) \end{aligned}$$