

t56_facirc_1 (TMSMipVBaicH- pipuDW2PBEcNu5Cetp9FhcK)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k6_margrel1 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_msafree2 : \iota \Rightarrow \iota$ be given. Let $k8_facirc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_facirc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $r1_circcomb : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_circcomb : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_msafree2 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_msualg_1 : \iota \Rightarrow o$ be given. Let $k2_circcomb : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v11_struct_0 : \iota \Rightarrow o$ be given. Let $v1_msualg_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 \\ & X1))) \Rightarrow (\forall X2. ((v1_relat_1 X2) \wedge ((v1_funct_1 X2) \wedge (v1_finseq_1 \\ & X2))) \Rightarrow (r1_circcomb (k5_circcomb X0 X1) (k5_circcomb X0 X2))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 \\ & X1))) \Rightarrow ((k2_msafree2 (k5_circcomb X0 X1) = k10_xtuple_0 X1) \wedge (k3_msafree2 \\ & (k5_circcomb X0 X1) = k1_tarski (k4_tarski X1 X0))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. k2_tarski X0 X1 = k2_xboole_0 (k1_tarski \\ & X0) (k1_tarski X1) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l1_msualg_1 X0)) \Rightarrow (\forall X1. \\ & ((\neg v2_struct_0 X1) \wedge (l1_msualg_1 X1)) \Rightarrow ((r1_circcomb X0 X1) \Rightarrow (\\ & (k3_msafree2 (k2_circcomb X0 X1) = k2_xboole_0 (k3_msafree2 X0) \\ & (k3_msafree2 X1)) \wedge (r1_tarski (k2_msafree2 (k2_circcomb X0 X1) \\ & (k2_xboole_0 (k2_msafree2 X0) (k2_msafree2 X1)))))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 \\ & X1))) \Rightarrow ((\neg v2_struct_0 (k5_circcomb X0 X1)) \wedge ((\neg v11_struct_0 (\\ & k5_circcomb X0 X1)) \wedge (v1_msualg_1 (k5_circcomb X0 X1)))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (v1_relat_1 (k10_finseq_1 X0 X1)) \wedge (v1_funct_1 \\ & (k10_finseq_1 X0 X1)) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. v1_finseq_1 (k10_finseq_1 X0 X1) \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 \\ & X1))) \Rightarrow ((\neg v11_struct_0 (k5_circcomb X0 X1)) \wedge ((v1_msualg_1 (k5_circcomb \\ & X0 X1)) \wedge (l1_msualg_1 (k5_circcomb X0 X1)))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((v1_funct_1 X3) \wedge \\ & ((v1_funct_2 X3 (k4_finseq_2 np_2 k6_margrel1) k6_margrel1) \wedge \\ & (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 (k4_finseq_2 np_2 \\ & k6_margrel1) k6_margrel1)))) \Rightarrow (k9_facirc_1 X0 X1 X2 X3 = k4_tarski \\ & (k10_finseq_1 (k4_tarski (k10_finseq_1 X0 X1) X3) X2) X3) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((v1_funct_1 X3) \wedge \\ & ((v1_funct_2 X3 (k4_finseq_2 np_2 k6_margrel1) k6_margrel1) \wedge \\ & (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 (k4_finseq_2 np_2 \\ & k6_margrel1) k6_margrel1)))) \Rightarrow (k8_facirc_1 X0 X1 X2 X3 = k2_circcomb \\ & (k5_circcomb X3 (k10_finseq_1 X0 X1)) (k5_circcomb X3 (k10_finseq_1 \\ & (k4_tarski (k10_finseq_1 X0 X1) X3) X2))) \end{aligned} \quad (10)$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((v1_funct_1 X3) \wedge \\ & ((v1_funct_2 X3 (k4_finseq_2 np_2 k6_margrel1) k6_margrel1) \wedge \\ & (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 (k4_finseq_2 np_2 \\ & k6_margrel1) k6_margrel1)))))) \Rightarrow (k3_msafree2 (k8_facirc_1 X0 \\ X1 X2 X3) = k2_tarski (k4_tarski (k10_finseq_1 X0 X1) X3) (k9_facirc_1 \\ X0 X1 X2 X3)) \end{aligned}$$