

# t56\_flang\_3 (TMGzpKwfTUS- bSd3htu3Q6eqywbNtY77pZWb)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k8\_afinsq\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_flang\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_flang\_3 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_flang\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k8\_flang\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_catalan2 : \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k8\_afinsq\_1 X0))) \Rightarrow (k2\_flang\_3 X0 X1 = k6\_flang\_1 X0 (k8\_flang\_1 X0 X1) X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k3\_catalan2 X0))) \Rightarrow (k2\_flang\_1 X0 \in k8\_flang\_1 X0 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k3\_catalan2 X0))) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k3\_catalan2 X0))) \Rightarrow ((k2\_flang\_1 X0 \in X1) \Rightarrow ((r1\_tarski X2 (k6\_flang\_1 X0 X2 X1)) \wedge (r1\_tarski X2 (k6\_flang\_1 X0 X1 X2)))))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k3\_catalan2 X0))) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k3\_catalan2 X0))) \Rightarrow ((k2\_flang\_1 X0 \in k6\_flang\_1 X0 X1 X2) \Leftrightarrow ((k2\_flang\_1 X0 \in X1) \wedge (k2\_flang\_1 X0 \in X2)))) \quad (4)$$

Assume the following.

$$\forall X0. k3\_catalan2 X0 = k8\_afinsq\_1 X0 \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k3\_catalan2 X0))) \Rightarrow (m1\_subset\_1 (k8\_flang\_1 X0 X1) (k1\_zfmisc\_1 (k3\_catalan2 X0))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(r1\_tarSKI X0 X1)\Leftrightarrow(\forall X2.(X2 \in X0)\Rightarrow (X2 \in X1)) \quad (7)$$

**Theorem 1**

$$\forall X0.\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k8\_afinsq\_1 X0)))\Rightarrow((k2\_flang\_1 X0 \in k2\_flang\_3 X0 X1)\Leftrightarrow(k2\_flang\_1 X0 \in X1))$$