

t56\_fomodel0  
(TMW2Lb9EHZ1GiAnmCkjocyrW8zckeg7q1cU)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $k5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X0) \wedge (v1\_relat\_1 X1)) \Rightarrow (v1\_relat\_1 (k2\_xboole\_0 X0 X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (v1\_relat\_1 X0) \Rightarrow (v1\_relat\_1 (k5\_relat\_1 X0 X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k2\_xboole\_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \vee (X3 \in X1))) \quad (3)$$

Assume the following.

$$\forall X0. (v1\_relat\_1 X0) \Leftrightarrow (\forall X1. \neg (X1 \in X0) \wedge (\forall X2. \forall X3. X1 \neq k4\_tarski X2 X3)) \quad (4)$$

Assume the following.

$$\forall X0. (v1\_relat\_1 X0) \Rightarrow (\forall X1. \forall X2. (v1\_relat\_1 X2) \Rightarrow ((X2 = k5\_relat\_1 X0 X1) \Leftrightarrow (\forall X3. \forall X4. (k4\_tarski X3 X4 \in X2) \Leftrightarrow ((X3 \in X1) \wedge (k4\_tarski X3 X4 \in X0))))) \quad (5)$$

**Theorem 1**

$$\forall X0. \forall X1. (v1\_relat\_1 X1) \Rightarrow (\forall X2. (v1\_relat\_1 X2) \Rightarrow (k5\_relat\_1 (k2\_xboole\_0 X1 X2) X0 = k2\_xboole\_0 (k5\_relat\_1 X1 X0) (k5\_relat\_1 X2 X0)))$$