

t56_group_2
(TMYjethaMB9Hv79dhU9zQ6MrjpV9nh9jfSs)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v2_group_1 : \iota \Rightarrow o$ be given. Let $v3_group_1 : \iota \Rightarrow o$ be given. Let $l3_algstr_0 : \iota \Rightarrow o$ be given. Let $m1_group_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u2_algstr_0 : \iota \Rightarrow \iota$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_realset1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((r1_tarski X0 X1) \wedge (r1_tarski X2 X3)) \Rightarrow (r1_tarski (k2_zfmisc_1 X0 X2) (k2_zfmisc_1 X1 X3)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((r1_tarski X0 X1) \Rightarrow ((k5_relat_1 (k5_relat_1 X2 X0) X1 = k5_relat_1 X2 X0) \wedge (k5_relat_1 (k5_relat_1 X2 X1) X0 = k5_relat_1 X2 X0))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((r1_tarski X0 X1) \wedge (r1_tarski X1 X2)) \Rightarrow (r1_tarski X0 X2) \quad (3)$$

Assume the following.

$$\forall X0.(l3_algstr_0 X0) \Rightarrow ((v1_funct_1 (u2_algstr_0 X0)) \wedge ((v1_funct_2 (u2_algstr_0 X0) (k2_zfmisc_1 (u1_struct_0 X0) (u1_struct_0 X0)) (u1_struct_0 X0)) \wedge (m1_subset_1 (u2_algstr_0 X0) (k1_zfmisc_1 (k2_zfmisc_1 (k2_zfmisc_1 (u1_struct_0 X0) (u1_struct_0 X0)) (u1_struct_0 X0)))))) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_group_1 X0) \wedge (l3_algstr_0 \\ X0))) \Rightarrow (\forall X1.((\neg v2_struct_0 X1) \wedge ((v2_group_1 X1) \wedge (l3_algstr_0 \\ X1))) \Rightarrow ((m1_group_2 X1 X0) \Leftrightarrow ((r1_tarski (u1_struct_0 X1) (u1_struct_0 \\ X0)) \wedge (u2_algstr_0 X1 = k1_realset1 (u2_algstr_0 X0) (u1_struct_0 \\ X1)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0) \Rightarrow (\forall X1.k1_realset1 X0 X1 = k5_relat_1 \\ X0 (k2_zfmisc_1 X1 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (7)$$

Theorem 1

$$\begin{aligned} \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_group_1 X0) \wedge ((v3_group_1 \\ X0) \wedge (l3_algstr_0 X0)))) \Rightarrow (\forall X1.((\neg v2_struct_0 X1) \wedge ((v2_group_1 \\ X1) \wedge ((v3_group_1 X1) \wedge (l3_algstr_0 X1)))) \Rightarrow (\forall X2.((\neg v2_struct_0 \\ X2) \wedge ((v2_group_1 X2) \wedge ((v3_group_1 X2) \wedge (l3_algstr_0 X2)))) \Rightarrow \\ (((m1_group_2 X0 X1) \wedge (m1_group_2 X1 X2)) \Rightarrow (m1_group_2 X0 X2)))) \end{aligned}$$