

# t56\_interval1

## (TMSQfBiqM5pfYwQ3SH5q53fNJzLegzBjitB)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v3\_roughs\_1 : \iota \Rightarrow o$  be given. Let  $l1\_orders\_2 : \iota \Rightarrow o$  be given. Let  $m2\_interval1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k14\_interval1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k16\_interval1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k4\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k15\_interval1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 X0))) \Rightarrow (k4\_subset\_1 X0 X1 X2 = k2\_xboole\_0 X1 X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. k1\_xtuple\_0 (k4\_tarski X0 X1) = X0 \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (((\neg v2\_struct\_0 X0) \wedge ((v3\_roughs\_1 X0) \wedge (l1\_orders\_2 X0))) \wedge ((m2\_interval1 X1 X0) \wedge (m2\_interval1 X2 X0))) \Rightarrow (m2\_interval1 (k16\_interval1 X0 X1 X2) X0) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (((\neg v2\_struct\_0 X0) \wedge ((v3\_roughs\_1 X0) \wedge (l1\_orders\_2 X0))) \wedge (m2\_interval1 X1 X0)) \Rightarrow (m1\_subset\_1 (k14\_interval1 X0 X1) (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \quad (4)$$

Assume the following.

$$\forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v3\_roughs\_1 X0) \wedge (l1\_orders\_2 X0))) \Rightarrow (\forall X1. (m2\_interval1 X1 X0) \Rightarrow (\forall X2. (m2\_interval1 X2 X0) \Rightarrow (k16\_interval1 X0 X1 X2 = k4\_tarski (k4\_subset\_1 (u1\_struct\_0 X0) (k14\_interval1 X0 X1) (k14\_interval1 X0 X2)) (k4\_subset\_1 (u1\_struct\_0 X0) (k15\_interval1 X0 X1) (k15\_interval1 X0 X2)))))) \quad (5)$$

Assume the following.

$$\forall X0.((\neg v2\_struct\_0 X0) \wedge ((v3\_roughs\_1 X0) \wedge (l1\_orders\_2 X0))) \Rightarrow (\forall X1.(m2\_interval1 X1 X0) \Rightarrow (k14\_interval1 X0 X1 = k1\_xtuple\_0 X1)) \quad (6)$$

**Theorem 1**

$$\forall X0.((\neg v2\_struct\_0 X0) \wedge ((v3\_roughs\_1 X0) \wedge (l1\_orders\_2 X0))) \Rightarrow (\forall X1.(m2\_interval1 X1 X0) \Rightarrow (\forall X2.(m2\_interval1 X2 X0) \Rightarrow (k14\_interval1 X0 (k16\_interval1 X0 X1 X2) = k4\_subset\_1 (u1\_struct\_0 X0) (k14\_interval1 X0 X1) (k14\_interval1 X0 X2))))$$