

t56_partfun1
 (TMJavEPnWXeDa1w1HVi8hJ59jMA2tpBStVf)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $r1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_partfun1 X0 X1) \Leftrightarrow (\exists X2. \\ ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \wedge (k2_xboole_0 X0 X1 = X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_xboole_0 (k9_xtuple_0 \\ X0) (k9_xtuple_0 X1)) \Rightarrow ((v1_relat_1 (k2_xboole_0 X0 X1)) \wedge (v1_funct_1 \\ (k2_xboole_0 X0 X1)))))) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_xboole_0 (k9_xtuple_0 \\ X0) (k9_xtuple_0 X1)) \Rightarrow (r1_partfun1 X0 X1))) \end{aligned}$$