

t57_bvfunc26 (TMGGX-
siU5PkcUTA3TPxKTzGCN2eSKgqbMWf)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_margrel1 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_bvfunc26 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_bvfunc_1 : \iota \Rightarrow \iota$ be given. Let $k1_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (r2_funct_2 X0 k6_margrel1 (\\ & k5_bvfunc_1 X0 X1 (k1_bvfunc_1 X0 X1)) (k12_bvfunc_1 X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge \\ & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X3.((v1_funct_1 X3) \wedge \\ & ((v1_funct_2 X3 X0 k6_margrel1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (r2_funct_2 X0 k6_margrel1 \\ & (k5_bvfunc_1 X0 X1 (k5_bvfunc26 X0 X2 X3)) (k5_bvfunc_1 X0 (k5_bvfunc_1 \\ & X0 X1 (k1_bvfunc_1 X0 X2)) (k1_bvfunc_1 X0 X3)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (r2_funct_2 X0 k6_margrel1 (\\ & k5_bvfunc_1 X0 X1 (k12_bvfunc_1 X0)) (k12_bvfunc_1 X0))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.(((v1_funct_1 X2)\wedge \\ & ((v1_funct_2 X2 X0 X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1)))))\wedge((v1_funct_1 X3)\wedge((v1_funct_2 X3 X0 X1)\wedge(m1_subset_1 \\ & X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))))\Rightarrow((r2_funct_2 X0 X1 X2 \\ & X3)\Leftrightarrow(X2 = X3)) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge(((v1_funct_1 \\ & X1)\wedge((v1_funct_2 X1 X0 k6_margrel1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))\wedge((v1_funct_1 X2)\wedge((v1_funct_2 \\ & X2 X0 k6_margrel1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k6_margrel1))))))\Rightarrow((v1_funct_1 (k5_bvfunc_1 X0 X1 X2))\wedge(\\ & (v1_funct_2 (k5_bvfunc_1 X0 X1 X2) X0 k6_margrel1)\wedge(m1_subset_1 \\ & (k5_bvfunc_1 X0 X1 X2) (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((v1_funct_1 X1)\wedge(\\ & (v1_funct_2 X1 X0 k6_margrel1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1))))))\Rightarrow((v1_funct_1 (k1_bvfunc_1 \\ & X0 X1))\wedge((v1_funct_2 (k1_bvfunc_1 X0 X1) X0 k6_margrel1)\wedge(m1_subset_1 \\ & (k1_bvfunc_1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow((v1_funct_1 (k12_bvfunc_1 X0))\wedge \\ & ((v1_funct_2 (k12_bvfunc_1 X0) X0 k6_margrel1)\wedge(m1_subset_1 \\ & (k12_bvfunc_1 X0) (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge(((v1_funct_1 \\ & X1)\wedge((v1_funct_2 X1 X0 k6_margrel1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))\wedge((v1_funct_1 X2)\wedge((v1_funct_2 \\ & X2 X0 k6_margrel1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k6_margrel1))))))\Rightarrow(k5_bvfunc_1 X0 X1 X2 = k5_bvfunc_1 X0 X2 \\ & X1)) \end{aligned} \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((v1_funct_1 X1)\wedge(\\ & (v1_funct_2 X1 X0 k6_margrel1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1))))))\Rightarrow(\forall X2.((v1_funct_1 X2)\wedge \\ & ((v1_funct_2 X2 X0 k6_margrel1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1))))))\Rightarrow(r2_funct_2 X0 k6_margrel1 \\ & (k5_bvfunc_1 X0 X1 (k5_bvfunc_26 X0 X1 X2)) (k12_bvfunc_1 X0))) \end{aligned}$$