

t57_bvfunc_1 (TMcKSjsQzGHYHW-
tYN62rCQAc3AgBP4CVbRd)

October 27, 2020

Let $v1_xboolean : \iota \Rightarrow o$ be given. Let $k7_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_margrel1 : \iota$ be given. Let $k3_xboolean : \iota \Rightarrow \iota$ be given. Let $k10_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboolean : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_margrel1 : \iota$ be given. Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (k10_xboolean k7_margrel1 X0 = X0) \quad (1)$$

Assume the following.

$$k7_margrel1 = k1_xboolean \quad (2)$$

Assume the following.

$$m1_subset_1 k7_margrel1 k6_margrel1 \quad (3)$$

Assume the following.

$$\forall X0.(v1_xboolean X0) \Rightarrow (\forall X1.(v1_xboolean X1) \Rightarrow (k7_xboolean X0 X1 = k3_xboolean (k10_xboolean X0 X1))) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xboolean X0) \wedge (v1_xboolean X1)) \Rightarrow (k7_xboolean X0 X1 = k7_xboolean X1 X0) \quad (5)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k6_margrel1) \Rightarrow (v1_xboolean X0) \quad (6)$$

Theorem 1

$$\forall X0.(v1_xboolean X0) \Rightarrow (k7_xboolean k7_margrel1 X0 = k3_xboolean X0)$$