

t57_group_11
(TMWYGkPxyinNtsq4h3eawkgUhcGWHa9c1zJ)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v2_group_1 : \iota \Rightarrow o$ be given. Let $v3_group_1 : \iota \Rightarrow o$ be given. Let $l3_algstr_0 : \iota \Rightarrow o$ be given. Let $m1_group_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_group_11 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k13_group_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k14_group_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k8_group_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_group_1 X0) \wedge ((v3_group_1 \\ & X0) \wedge (l3_algstr_0 X0)))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 \\ & X0)) \Rightarrow (\forall X2.(m1_group_2 X2 X0) \Rightarrow (\forall X3.(m1_group_2 \\ & X3 X0) \Rightarrow ((m1_group_2 X2 X3) \Rightarrow ((r1_tarski (k13_group_2 X0 X2 X1) (\\ & k13_group_2 X0 X3 X1)) \wedge (r1_tarski (k14_group_2 X0 X2 X1) (k14_group_2 \\ & X0 X3 X1))))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1_tarski X0 X1) \wedge (r1_xboole_0 X1 X2)) \Rightarrow (r1_xboole_0 X0 X2) \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_group_1 X0) \wedge ((v3_group_1 \\ & X0) \wedge (l3_algstr_0 X0)))) \Rightarrow (\forall X1.(m1_group_2 X1 X0) \Rightarrow (k8_group_2 \\ & X0 X1 = u1_struct_0 X1)) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_group_1 X0) \wedge ((v3_group_1 \\ & X0) \wedge (l3_algstr_0 X0)))) \Rightarrow (\forall X1.(m1_group_2 X1 X0) \Rightarrow (\forall X2. \\ & (m1_group_2 X2 X0) \Rightarrow (k4_group_11 X0 X1 X2 = ReplSep (toset (\lambda X3 : \\ & \iota.m1_subset_1 X3 (u1_struct_0 X0))) (\lambda X3 : \iota.\neg r1_xboole_0 \\ & (k13_group_2 X0 X2 X3) (k8_group_2 X0 X1)) (\lambda X3 : \iota.X3)))) \end{aligned} \tag{4}$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarSKI X0 X1)\Leftrightarrow(\forall X2.(X2 \in X0)\Rightarrow (X2 \in X1)) \quad (5)$$

Theorem 1

$$\begin{aligned} &\forall X0.((\neg v2_struct_0 X0)\wedge((v2_group_1 X0)\wedge((v3_group_1 \\ &X0)\wedge(l3_algstr_0 X0))))\Rightarrow(\forall X1.(m1_group_2 X1 X0)\Rightarrow(\forall X2. \\ &(m1_group_2 X2 X0)\Rightarrow(\forall X3.(m1_group_2 X3 X0)\Rightarrow((m1_group_2 \\ &X2 X3)\Rightarrow(r1_tarSKI (k4_group_11 X0 X1 X2) (k4_group_11 X0 X1 X3)))))) \end{aligned}$$