

t57_relat_1 (TMRyPHNxrvt- pReBx3UMZpQMCawwwgFN6Dz)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X0) \Rightarrow (v1_relat_1 (k5_relat_1 X0 X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k9_xtuple_0 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. k4_tarski X2 X3 \in X0)) \quad (2)$$

Assume the following.

$$\forall X0. (v1_relat_1 X0) \Rightarrow (\forall X1. \forall X2. (v1_relat_1 X2) \Rightarrow ((X2 = k5_relat_1 X0 X1) \Leftrightarrow (\forall X3. \forall X4. (k4_tarski X3 X4 \in X2) \Leftrightarrow ((X3 \in X1) \wedge (k4_tarski X3 X4 \in X0))))) \quad (3)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. (v1_relat_1 X2) \Rightarrow ((X0 \in k9_xtuple_0 (k5_relat_1 X2 X1)) \Leftrightarrow ((X0 \in X1) \wedge (X0 \in k9_xtuple_0 X2)))$$