

# t57\_xboolean

## (TMboLk3rCHUqWirBctTg5Z1eftuvdBzCewj)

October 27, 2020

Let  $v1\_xboolean : \iota \Rightarrow o$  be given. Let  $k8\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_xboolean : \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xboolean X0) \Rightarrow (\forall X1.(v1\_xboolean X1) \Rightarrow (\forall X2. \\ & (v1\_xboolean X2) \Rightarrow (k5\_xboolean X0 (k4\_xboolean X1 X2) = k4\_xboolean \\ & (k5\_xboolean X0 X1) (k5\_xboolean X0 X2)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xboolean X0) \Rightarrow (\forall X1.(v1\_xboolean X1) \Rightarrow (\forall X2. \\ & (v1\_xboolean X2) \Rightarrow (k8\_xboolean X0 (k8\_xboolean X1 X2) = k4\_xboolean \\ & (k5\_xboolean (k3\_xboolean X0) X1) (k5\_xboolean (k3\_xboolean X0) \\ & X2)))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0.(v1\_xboolean X0) \Rightarrow (\forall X1.(v1\_xboolean X1) \Rightarrow (k6\_xboolean X0 (k4\_xboolean X0 X1) = k6\_xboolean X0 X1)) \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xboolean X0) \wedge (v1\_xboolean X1)) \Rightarrow (v1\_xboolean (k4\_xboolean X0 X1)) \tag{4}$$

Assume the following.

$$\forall X0.(v1\_xboolean X0) \Rightarrow (v1\_xboolean (k3\_xboolean X0)) \tag{5}$$

Assume the following.

$$\forall X0.(v1\_xboolean X0) \Rightarrow (\forall X1.(v1\_xboolean X1) \Rightarrow (k6\_xboolean X0 X1 = k5\_xboolean (k3\_xboolean X0) X1)) \tag{6}$$

### Theorem 1

$$\forall X0.(v1\_xboolean X0) \Rightarrow (\forall X1.(v1\_xboolean X1) \Rightarrow (k8\_xboolean X0 (k8\_xboolean X0 X1) = k6\_xboolean X0 X1))$$