

## t57\_xxreal\_1

(TMQgNaSs1nYht9xjXmxGLrtREhjtaB74trW)

October 27, 2020

Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xxreal\_0 X2) \Rightarrow (\forall X3.(v1\_xxreal\_0 X3) \Rightarrow (((r1\_xxreal\_0 \\ & X0 X1) \wedge (r1\_tarski (k1\_xxreal\_1 X0 X1) (k2\_xxreal\_1 X2 X3))) \Rightarrow (( \\ & r1\_xxreal\_0 X2 X0) \wedge (\neg r1\_xxreal\_0 X3 X1)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xxreal\_0 X2) \Rightarrow (\forall X3.(v1\_xxreal\_0 X3) \Rightarrow ((r1\_tarski ( \\ & k3\_xxreal\_1 X0 X1) (k1\_xxreal\_1 X2 X3)) \Rightarrow ((r1\_xxreal\_0 X1 X0) \vee ( \\ & r1\_xxreal\_0 X2 X0) \wedge (r1\_xxreal\_0 X1 X3)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xxreal\_0 X2) \Rightarrow (\forall X3.(v1\_xxreal\_0 X3) \Rightarrow ((r1\_xxreal\_0 \\ & X2 X3) \Rightarrow ((r1\_xxreal\_0 X1 X0) \vee (r1\_tarski (k1\_xxreal\_1 X1 X2) (k3\_xxreal\_1 \\ & X0 X3)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xxreal\_0 X2) \Rightarrow (((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 X1 X2)) \Rightarrow \\ & (r1\_xxreal\_0 X0 X2)))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (r1\_tarski \\ & (k2\_xxreal\_1 X0 X1) (k1\_xxreal\_1 X0 X1))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((r1\_tarSKI X0 X1)\wedge(r1\_tarSKI X1 X2))\Rightarrow(r1\_tarSKI X0 X2) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xxreal\_0 X0)\wedge(v1\_xxreal\_0 X1))\Rightarrow(r1\_xxreal\_0 X0 X0) \quad (7)$$

**Theorem 1**

$$\forall X0.(v1\_xxreal\_0 X0)\Rightarrow(\forall X1.(v1\_xxreal\_0 X1)\Rightarrow(\forall X2.(v1\_xxreal\_0 X2)\Rightarrow(\forall X3.(v1\_xxreal\_0 X3)\Rightarrow((r1\_tarSKI (k3\_xxreal\_1 X0 X1) (k2\_xxreal\_1 X2 X3))\Rightarrow((r1\_xxreal\_0 X1 X0)\vee(r1\_xxreal\_0 X2 X0)\wedge(\neg r1\_xxreal\_0 X3 X1)))))))$$