

t58_complfld (TMPGFk-
JeH8DtHGCvtEa3v3eG7ybbTkXMZAv)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_complfld : \iota$ be given. Let $k17_complex1 : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k4_struct_0 : \iota \Rightarrow \iota$ be given. Let $k5_complex1 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Assume the following.

$$k4_struct_0 \ k1_complfld = k5_complex1 \tag{1}$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 \ X0) \Rightarrow ((k17_complex1 \ X0 = k6_numbers) \Rightarrow (X0 = k6_numbers)) \tag{2}$$

Assume the following.

$$k6_numbers = k1_xboole_0 \tag{3}$$

Assume the following.

$$k5_complex1 = k1_xboole_0 \tag{4}$$

Assume the following.

$$\forall X0.(m1_subset_1 \ X0 \ (u1_struct_0 \ k1_complfld)) \Rightarrow (v1_xcmplx_0 \ X0) \tag{5}$$

Theorem 1

$$\forall X0.(m1_subset_1 \ X0 \ (u1_struct_0 \ k1_complfld)) \Rightarrow ((k17_complex1 \ X0 = k6_numbers) \Rightarrow (X0 = k4_struct_0 \ k1_complfld))$$