

t58_eqrel_1

(TMXG6HNuuPVjyx2ZV4kSNYHRwju6Fdfwwzr)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_binop_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k16_funct_3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_partfun1 : \iota \Rightarrow \iota$ be given. Let $k1_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_relat_1 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(\neg v1_xboole_0 X1) \Rightarrow \\
 & (\forall X2.(\neg v1_xboole_0 X2) \Rightarrow (\forall X3.(\neg v1_xboole_0 X3) \Rightarrow \\
 & (\forall X4.((v1_funct_1 X4) \wedge ((v1_funct_2 X4 X0 X1) \wedge (m1_subset_1 \\
 & X4 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \Rightarrow (\forall X5.((v1_funct_1 \\
 & X5) \wedge ((v1_funct_2 X5 X2 X3) \wedge (m1_subset_1 X5 (k1_zfmisc_1 (k2_zfmisc_1 \\
 & X2 X3)))))) \Rightarrow (\forall X6.(m1_subset_1 X6 X0) \Rightarrow (\forall X7.(m1_subset_1 \\
 & X7 X2) \Rightarrow (k2_binop_1 X0 X2 (k2_zfmisc_1 X1 X3) (k16_funct_3 X0 X2 X1 \\
 & X3 X4 X5) X6 X7 = k4_tarski (k3_funct_2 X0 X1 X4 X6) (k3_funct_2 X2 X3 \\
 & X5 X7))))))))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.\forall X1.(m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.(X0 \in X1) \Rightarrow (k1_funct_1 (k4_relat_1 X1) X0 = X0) \tag{3}$$

Assume the following.

$$\forall X0.k6_partfun1 X0 = k4_relat_1 X0 \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0)\wedge \\ & (((v1_funct_1 X2)\wedge((v1_funct_2 X2 X0 X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))))))\wedge(m1_subset_1 X3 X0)))\Rightarrow(k3_funct_2 X0 \\ & X1 X2 X3 = k1_funct_1 X2 X3) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0)\wedge \\ & ((\neg v1_xboole_0 X1)\wedge((m1_subset_1 X2 X0)\wedge(m1_subset_1 X3 X1))))\Rightarrow \\ & (k1_domain_1 X0 X1 X2 X3 = k4_tarski X2 X3) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_relat_1 (k4_relat_1 X0))\wedge((v4_relat_1 (k4_relat_1 \\ & X0) X0)\wedge((v1_funct_1 (k4_relat_1 X0))\wedge(v1_partfun1 (k4_relat_1 \\ & X0) X0))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_partfun1 (k6_partfun1 X0) X0)\wedge(m1_subset_1 (k6_partfun1 \\ & X0) (k1_zfmisc_1 (k2_zfmisc_1 X0 X0))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0)\wedge \\ & (((v1_funct_1 X2)\wedge((v1_funct_2 X2 X0 X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))))))\wedge(m1_subset_1 X3 X0)))\Rightarrow(m1_subset_1 (\\ & k3_funct_2 X0 X1 X2 X3) X1) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1)))\Rightarrow((v1_partfun1 X2 X0)\Rightarrow(v1_funct_2 X2 X0 X1)) \end{aligned} \quad (10)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.(\neg v1_xboole_0 X1)\Rightarrow \\ & (\forall X2.(\neg v1_xboole_0 X2)\Rightarrow(\forall X3.((v1_funct_1 X3)\wedge \\ & ((v1_funct_2 X3 X0 X1)\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1))))))\Rightarrow(\forall X4.(m1_subset_1 X4 X0)\Rightarrow(\forall X5.(m1_subset_1 \\ & X5 X2)\Rightarrow(k2_binop_1 X0 X2 (k2_zfmisc_1 X1 X2) (k16_funct_3 X0 X2 X1 \\ & X2 X3 (k6_partfun1 X2)) X4 X5 = k1_domain_1 X1 X2 (k3_funct_2 X0 X1 \\ & X3 X4) X5)))))) \end{aligned}$$