

t58_finseq_1 (TMJD-
VDzHtn4zV1VFb9oU3M3aiELZwmUvXDK)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k16_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_card_1 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow ((r1_tarski (k9_xtuple_0 X1) X0) \Rightarrow (k5_relat_1 X1 X0 = X1)) \quad (1)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Leftrightarrow (r1_tarski (k2_finseq_1 X0) (k2_finseq_1 X1)))) \quad (2)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k3_finseq_1 X0 = k1_card_1 X0) \quad (3)$$

Assume the following.

$$\forall X0. (v1_finset_1 X0) \Rightarrow ((v1_finset_1 (k1_card_1 X0)) \wedge (v1_card_1 (k1_card_1 X0))) \quad (4)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (m2_subset_1 (k3_finseq_1 X0) k1_numbers k5_numbers) \quad (5)$$

Assume the following.

$$\forall X0. v1_card_1 (k1_card_1 X0) \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow ((X1 = k3_finseq_1 \\ & X0) \Leftrightarrow (k2_finseq_1 X1 = k9_xtuple_0 X0))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge ((\\ & v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow (k16_finseq_1 X0 X1 = k5_relat_1 \\ & X1 (k2_finseq_1 X0))) \end{aligned} \quad (8)$$

Assume the following.

$$\forall X0.((v3_ordinal1 X0) \wedge (v1_finset_1 X0)) \Rightarrow (v7_ordinal1 X0) \quad (9)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finset_1 X0))) \end{aligned} \quad (10)$$

Assume the following.

$$\forall X0.(v1_card_1 X0) \Rightarrow (v3_ordinal1 X0) \quad (11)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge ((\\ & v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow ((r1_xxreal_0 (k3_finseq_1 \\ & X1) X0) \Rightarrow (k16_finseq_1 X0 X1 = X1))) \end{aligned}$$