

# t58\_interva1 (TMNmQL- CmhS3w2KXbD45aqNbdKuvFfLwJrqc)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v3\_roughs\_1 : \iota \Rightarrow o$  be given. Let  $l1\_orders\_2 : \iota \Rightarrow o$  be given. Let  $m2\_interval1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k14\_interval1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k17\_interval1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k4\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k15\_interval1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. k1\_xtuple\_0 (k4\_tarski X0 X1) = X0 \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (((\neg v2\_struct\_0 X0) \wedge ((v3\_roughs\_1 X0) \wedge (l1\_orders\_2 X0) \wedge (l1\_orders\_2 X0))) \wedge ((m2\_interval1 X1 X0) \wedge (m2\_interval1 X2 X0))) \Rightarrow (m2\_interval1 (k17\_interval1 X0 X1 X2) X0) \quad (2)$$

Assume the following.

$$\forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v3\_roughs\_1 X0) \wedge (l1\_orders\_2 X0))) \Rightarrow (\forall X1. (m2\_interval1 X1 X0) \Rightarrow (\forall X2. (m2\_interval1 X2 X0) \Rightarrow (k17\_interval1 X0 X1 X2 = k4\_tarski (k9\_subset\_1 (u1\_struct\_0 X0) (k14\_interval1 X0 X1) (k14\_interval1 X0 X2)) (k9\_subset\_1 (u1\_struct\_0 X0) (k15\_interval1 X0 X1) (k15\_interval1 X0 X2)))))) \quad (3)$$

Assume the following.

$$\forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v3\_roughs\_1 X0) \wedge (l1\_orders\_2 X0))) \Rightarrow (\forall X1. (m2\_interval1 X1 X0) \Rightarrow (k14\_interval1 X0 X1 = k1\_xtuple\_0 X1)) \quad (4)$$

## Theorem 1

$$\forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v3\_roughs\_1 X0) \wedge (l1\_orders\_2 X0))) \Rightarrow (\forall X1. (m2\_interval1 X1 X0) \Rightarrow (\forall X2. (m2\_interval1 X2 X0) \Rightarrow (k14\_interval1 X0 (k17\_interval1 X0 X1 X2) = k9\_subset\_1 (u1\_struct\_0 X0) (k14\_interval1 X0 X1) (k14\_interval1 X0 X2))))$$