

t58_moebius1

(TMNw2WYkwJxf9qanP8HE3mrKPrnwgo3pviy)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_moebius1 : \iota \Rightarrow \iota$ be given. Let $r1_nat_d : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((\\ r1_nat_d X1 X0) \Rightarrow ((r1_xxreal_0 X0 k6_numbers) \vee (r1_xxreal_0 X1 \\ X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 \\ X0 X1) \Rightarrow ((v1_xboole_0 X0) \vee ((v2_xxreal_0 X1) \vee (v3_xxreal_0 X0)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (3)$$

Assume the following.

$$\forall X0.((\neg v1_xboole_0 X0) \wedge (v7_ordinal1 X0)) \Rightarrow (r1_nat_d (\\ k6_moebius1 X0) X0) \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (5)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (6)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (7)$$

Assume the following.

$$\exists X0.(v1_xboole_0 X0) \wedge ((v1_xcmplx_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (v1_xreal_0 X0))) \quad (8)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (9)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (10)$$

Assume the following.

$$\forall X0.((\neg v1_xboole_0 X0) \wedge (v7_ordinal1 X0)) \Rightarrow (m2_subset_1 (k6_moebius1 X0) k1_numbers k5_numbers) \quad (11)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (12)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (13)$$

Assume the following.

$$\forall X0.((v1_xxreal_0 X0) \wedge (v2_xxreal_0 X0)) \Rightarrow ((\neg v1_xboole_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (\neg v3_xxreal_0 X0))) \quad (14)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow ((v7_ordinal1 X0) \wedge (\neg v3_xxreal_0 X0)) \quad (15)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_xreal_0 X0) \quad (16)$$

Theorem 1

$$\forall X0.((\neg v1_xboole_0 X0) \wedge (v7_ordinal1 X0)) \Rightarrow (r1_xxreal_0 (k6_moebius1 X0) X0)$$