

t58_sprect_2

(TMaa8PYyySXVEw1bjCtcEb7mGnzMDJdHf45)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v3_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_6 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $v1_topreal1 : \iota \Rightarrow o$ be given. Let $v2_topreal1 : \iota \Rightarrow o$ be given. Let $v1_goboard5 : \iota \Rightarrow o$ be given. Let $v2_goboard5 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k18_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k3_topreal1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k19_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v1_xboole_0 X0) \wedge (\neg v3_funct_1 X0) \wedge ((v1_finseq_6 \\ & X0 (u1_struct_0 (k15_euclid np_2))) \wedge ((v1_topreal1 X0) \wedge ((v2_topreal1 \\ & X0) \wedge ((v1_goboard5 X0) \wedge ((v2_goboard5 X0) \wedge (m2_finseq_1 X0 (u1_struct_0 \\ & (k15_euclid np_2)))))))))) \Rightarrow (\neg r1_xxreal_0 (k18_euclid (k19_pscomp_1 \\ & (k3_topreal1 np_2 X0))) (k18_euclid (k18_pscomp_1 (k3_topreal1 \\ & np_2 X0)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. ((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (r1_xxreal_0 X0 X0) \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \tag{4}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(m1_finseq_1 X1 (u1_struct_0 (k15_euclid X0))))\Rightarrow(m1_subset_1 (k3_topreal1 X0 X1) (k1_zfmisc_1 (u1_struct_0 (k15_euclid X0)))) \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2))))\Rightarrow(m1_subset_1 (k18_pscomp_1 X0) (u1_struct_0 (k15_euclid np_2))) \quad (7)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2)))\Rightarrow(m1_subset_1 (k18_euclid X0) k1_numbers) \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xxreal_0 X0) \quad (10)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (11)$$

Theorem 1

$$\forall X0.((\neg v1_xboole_0 X0)\wedge((\neg v3_funct_1 X0)\wedge((v1_finseq_6 X0 (u1_struct_0 (k15_euclid np_2)))\wedge((v1_topreal1 X0)\wedge((v2_topreal1 X0)\wedge((v1_goboard5 X0)\wedge((v2_goboard5 X0)\wedge(m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2))))))))))\Rightarrow(k18_pscomp_1 (k3_topreal1 np_2 X0)\neq k19_pscomp_1 (k3_topreal1 np_2 X0))$$