

## t59\_int\_1

(TMHN8rj7jGmFBiUqh3v6ymheoFz2FfdNV87)

October 27, 2020

Let  $v1\_int\_1 : \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_int\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_int\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $np\_1 : \iota$  be given. Let  $k6\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_xcmplx\_0 : \iota \Rightarrow \iota$  be given. Let  $v2\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_0 : \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (X0 = k1\_xboole\_0) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (k3\_xcmplx\_0 np\_1 X0 = X0) \quad (2)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (k3\_xcmplx\_0 X0 k6\_numbers = k6\_numbers) \quad (3)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (k2\_xcmplx\_0 X0 k6\_numbers = X0) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xcmplx\_0 X0) \wedge (v1\_xcmplx\_0 X1)) \Rightarrow (k6\_xcmplx\_0 (k4\_xcmplx\_0 X0) (k4\_xcmplx\_0 X1) = k6\_xcmplx\_0 X1 X0) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1\_xcmplx\_0 X0) \wedge ((v1\_xcmplx\_0 X1) \wedge (v1\_xcmplx\_0 X2))) \Rightarrow (k2\_xcmplx\_0 (k2\_xcmplx\_0 X0 X1) X2 = k2\_xcmplx\_0 X0 (k2\_xcmplx\_0 X1 X2)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1\_xcmplx\_0 X0)\wedge((v1\_xcmplx\_0 X1)\wedge(v1\_xcmplx\_0 X2)))\Rightarrow(k3\_xcmplx\_0 (k2\_xcmplx\_0 X0 X1) X2 = k2\_xcmplx\_0 (k3\_xcmplx\_0 X0 X2) (k3\_xcmplx\_0 X1 X2)) \quad (7)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0)\Rightarrow(k3\_xcmplx\_0 X0 (k4\_xcmplx\_0 np\_1) = k4\_xcmplx\_0 X0) \quad (8)$$

Assume the following.

$$((v2\_xreal\_0 np\_1)\wedge(m2\_subset\_1 np\_1 k1\_numbers k5\_numbers))\wedge((m1\_subset\_1 np\_1 k5\_numbers)\wedge(m1\_subset\_1 np\_1 k1\_numbers)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xcmplx\_0 X0)\wedge(v1\_xcmplx\_0 X1))\Rightarrow(k2\_xcmplx\_0 X0 (k4\_xcmplx\_0 X1) = k6\_xcmplx\_0 X0 X1) \quad (10)$$

Assume the following.

$$(m2\_subset\_1 np\_0 k1\_numbers k5\_numbers)\wedge((m1\_subset\_1 np\_0 k5\_numbers)\wedge(m1\_subset\_1 np\_0 k1\_numbers)) \quad (11)$$

Assume the following.

$$v1\_xboole\_0 np\_0 \quad (12)$$

Assume the following.

$$k6\_xcmplx\_0 np\_0 np\_1 = k4\_xcmplx\_0 np\_1 \quad (13)$$

Assume the following.

$$k2\_xcmplx\_0 (k4\_xcmplx\_0 np\_1) np\_1 = np\_0 \quad (14)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (15)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0)\Rightarrow(k4\_xcmplx\_0 (k4\_xcmplx\_0 X0) = X0) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0)\wedge(v1\_xreal\_0 X1))\Rightarrow(v1\_xreal\_0 (k6\_xcmplx\_0 X0 X1)) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xcmplx\_0 X0)\wedge(v1\_xcmplx\_0 X1))\Rightarrow(v1\_xcmplx\_0 (k2\_xcmplx\_0 X0 X1)) \quad (18)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_int\_1 X0)\wedge(v1\_int\_1 X1))\Rightarrow(v1\_int\_1 (k3\_xcmplx\_0 X0 X1)) \quad (19)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_int\_1 X0)\wedge(v1\_int\_1 X1))\Rightarrow(v1\_int\_1 (k5\_int\_1 X0 X1)) \quad (20)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0)\Rightarrow(v1\_xcmplx\_0 (k4\_xcmplx\_0 X0)) \quad (21)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1\_int\_1 X0)\Rightarrow(\forall X1.(v1\_int\_1 X1)\Rightarrow(((X1\neq k6\_numbers)\Rightarrow \\ (k6\_int\_1 X0 X1 = k6\_xcmplx\_0 X0 (k3\_xcmplx\_0 (k5\_int\_1 X0 X1) X1)))\wedge \\ ((X1 = k6\_numbers)\Rightarrow(k6\_int\_1 X0 X1 = k6\_numbers)))) \end{aligned} \quad (22)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xcmplx\_0 X0)\wedge(v1\_xcmplx\_0 X1))\Rightarrow(k3\_xcmplx\_0 X0 X1 = k3\_xcmplx\_0 X1 X0) \quad (23)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xcmplx\_0 X0)\wedge(v1\_xcmplx\_0 X1))\Rightarrow(k2\_xcmplx\_0 X0 X1 = k2\_xcmplx\_0 X1 X0) \quad (24)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0)\Rightarrow(v1\_xcmplx\_0 X0) \quad (25)$$

Assume the following.

$$\forall X0.(v1\_int\_1 X0)\Rightarrow(v1\_xreal\_0 X0) \quad (26)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers)\Rightarrow(v1\_xcmplx\_0 X0) \quad (27)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers)\Rightarrow(v1\_xreal\_0 X0) \quad (28)$$

**Theorem 1**

$$\begin{aligned} \forall X0.(v1\_int\_1 X0)\Rightarrow(\forall X1.(v1\_int\_1 X1)\Rightarrow((X0\neq k6\_numbers)\Rightarrow \\ (X1 = k2\_xcmplx\_0 (k3\_xcmplx\_0 (k5\_int\_1 X1 X0) X0) (k6\_int\_1 X1 \\ X0)))) \end{aligned}$$